



6/7/75



09/889326

JG18 Rec'd PCT/PTO 1 1 JUL 2001

**[001] METHOD OF ELECTROCHEMICALLY DETECTING NUCLEIC ACID
OLIGOMER HYBRIDIZATION EVENTS**

[002] FIELD OF THE INVENTION

[003] The present invention is directed to a modified nucleic acid oligomer, as well as a method of electrochemically detecting sequence-specific nucleic acid oligomer hybridization events.

[004] BACKGROUND OF THE INVENTION

[005] Generally, gel-electrophoretic methods with autoradiographical or optical detection are used for DNA and RNA sequence analysis, for example in disease diagnosis, toxicological test procedures, genetic research and development, and the agricultural and pharmaceutical sectors.

[006] In the most significant gel-electrophoretic method with optical detection, the Sanger method, a solution containing DNA is divided into four samples. To differentiate the four samples, the primer (complementary starting sequence for replication) of each sample is covalently modified with a fluorescent dye that emits at a distinct wavelength. Starting at the primer, each sample is enzymatically replicated by DNA polymerase I. In addition to the requisite deoxyribonucleoside triphosphates of the bases A (adenine), T (thymine), C (cytosine), and G (guanine), each reaction mixture also contains sufficient 2',3'-dideoxy analog of one of these nucleoside triphosphates as a blocking base (one of each of the four possible blocking bases per sample) in order to terminate replication at all possible binding sites. After the four samples are combined, all lengths of replicated DNA fragments having blocking-base-specific fluorescence result and can be sorted gel-electrophoretically by length and characterized by fluorescent spectroscopy.

[007] Another optical detection method is based on the attachment of fluorescent dyes such as ethidium bromide on oligonucleotides. In comparison with a free solution of the dye, the fluorescence of such dyes changes

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
210

drastically upon association with double-stranded DNA or RNA and can therefore be used to detect hybridized DNA or RNA.

[008] In radio labeling, ^{32}P is built into the phosphate skeleton of the oligonucleotides, ^{32}P usually being added to the 5'-hydroxyl end by polynucleotide kinase. Thereafter, the labeled DNA is preferably cleaved, under defined conditions, at one of each of the four nucleotide types, such that an average of one cleavage per chain results. Thus, for a given base type, there are present in the reaction mixture chains extending from the ^{32}P -label to the position of that base (if there are multiple appearances of the base, chains of varying lengths will result accordingly). The four fragment mixtures are then gel-electrophoretically separated on four lanes. Thereafter, an autoradiogram of the gel is prepared, from which the sequence can be read directly.

[009] Some years ago, a further method of DNA sequencing was developed on the basis of optical (or autoradiographical) detection, namely sequencing by oligomer hybridization (cf. e.g. Drmanac et al., Genomics 4, (1989), pp. 114-128 or Bains et al., Theor. Biol. 135, (1988), pp. 303-307). In this method, a complete set of short oligonucleotides or nucleic acid oligomers (probe oligonucleotides), e.g. all 65,536 possible combinations of bases A, T, C, and G of an oligonucleotide octamer, are bound to a support material. The attachment occurs in an ordered grid comprising 65,536 test sites, each rather large amount of an oligonucleotide combination defining one test site, and the position of each individual test site (oligonucleotide combination) being known. On such a hybridization matrix, the oligomer chip, a DNA fragment whose sequence is to be determined (the target) is labeled with fluorescent dye (or ^{32}P) and hybridized under conditions that allow only one specific double-strand formation. In this way, the target DNA fragment binds only to those nucleic acid oligomers (in this example to the octamers) whose complementary sequence corresponds exactly to a portion (an octamer) of its own sequence. Thus, all of the nucleic acid oligomer sequences (octamer sequences) present in the fragment are determined by optical (or autoradiographical) detection of

the binding position of the hybridized DNA fragment. Due to the overlapping of neighboring nucleic acid oligomer sequences, the sequential sequence of the DNA fragment can be determined using appropriate mathematical algorithms. One of the advantages of this method lies in the miniaturization of the sequencing and thus in the enormous amount of data that can be simultaneously captured in one operation. In addition, primer and gel-electrophoretic separation of the DNA fragments can be dispensed with. This principle is exemplified in Figure 1 for a 13-base-long DNA fragment.

[010] The use of radioactive labels in DNA/RNA sequencing is associated with several disadvantages, such as elaborate, legally required safety precautions in dealing with radioactive materials, radiation exposure, limited spatial resolution capacity (maximum 1 mm²), and sensitivity that is high only when the radiation of the radioactive fragments act on an X-ray film for an appropriately long time (hours to days). Although the spatial resolution can be increased by additional hardware and software, and the detection time can be decreased by using beta scanners, both of these involve considerable additional costs.

[011] Some of the fluorescent dyes that are commonly used to label the DNA (e.g. ethidium bromide) are mutagenic and require appropriate safety precautions, as does the use of autoradiography. In nearly every case, the use of optical detection requires the use of one or more laser systems, and thus experienced personnel and appropriate safety precautions. The actual detection of the fluorescence requires additional hardware, such as optical components for amplification and, in the case of varying excitation and query wavelengths as in the Sanger method, a control system. Thus, depending on the required excitation wavelengths and the desired detection performance, considerable investment costs may result. In sequencing by hybridization on an oligomer chip, detection is even more costly since, in addition to the excitation system, high-resolution CCD cameras (charge coupled device cameras) are needed for the two-dimensional detection of fluorescent spots.

[012] Thus, although quantitative and extremely sensitive methods for DNA/RNA sequencing exist, these methods are time consuming, require painstaking sample preparation and expensive equipment, and are generally not available as portable systems.

[013] DESCRIPTION OF THE INVENTION

[014] Therefore, it is the object of the present invention to create for detecting nucleic acid oligomer hybrids an apparatus and a method that do not exhibit the disadvantages of the background art.

[015] According to the present invention, this object is solved by the modified nucleic acid oligomer according to independent claim 1, the method of producing a modified nucleic acid oligomer according to independent claim 21, the modified conductive surface according to independent claim 29, the method of producing a modified conductive surface according to independent claim 44, and a method of electrochemically detecting nucleic acid oligomer hybridization events according to independent claim 48.

[016] The following abbreviations and terms will be used in the context of the present invention:

[017]

Genetics

DNA	deoxyribonucleic acid
RNA	ribonucleic acid
PNA	peptide nucleic acid (synthetic DNA or RNA in which the sugar-phosphate moiety is replaced by an amino acid. If the sugar-phosphate moiety is replaced by the -NH-(CH ₂) ₂ -N(COCH ₂ -base)-CH ₂ CO- moiety, PNA will hybridize with DNA.)
A	adenine
G	guanine
C	cytosine

T	thymine
U	uracil
base	A, G, T, C, or U
bp	base pair
nucleic acid	At least two covalently-joined nucleotides or at least two covalently-joined pyrimidine (e.g. cytosine, thymine, or uracil) or purine bases (e.g. adenine or guanine). The term nucleic acid refers to any backbone of the covalently-linked pyrimidine or purine bases, such as the sugar-phosphate backbone of DNA, cDNA, or RNA, a peptide backbone of PNA, or analogous structures (e.g. a phosphoramidate, thiophosphate, or dithiophosphate backbone). The essential feature of a nucleic acid according to the present invention is that it can sequence-specifically bind naturally occurring cDNA or RNA.
nucleic acid oligomer	Nucleic acid of a base length that is not further specified (e.g. nucleic acid octamer: a nucleic acid having any backbone in which 8 pyrimidine or purine bases are covalently bound to one another).
oligomer	Equivalent to nucleic acid oligomer.
oligonucleotide	Equivalent to oligomer or nucleic acid oligomer, e.g. a DNA, PNA, or RNA fragment of a base length that is not further specified.
oligo	Abbreviation for oligonucleotide.
primer	Initial complementary fragment of an oligonucleotide, the base length of the primer being only approx. 4-8 bases. Serves as the starting point for enzymatic replication of the oligonucleotide.
mismatch	To form the Watson-Crick double-stranded oligonucleotide structure, the two single-strands hybridize in such a way that the A (or C) base of one strand forms hydrogen bonds with the

	T (or G) base of the other strand (in RNA, T is replaced by uracil). Any other base pairing does not form hydrogen bonds, distorts the structure and is referred to as a "mismatch."
ss	single-strand
ds	double-strand

[018]

Photoinducibly and Chemically-Inducibly Redox-Active Moieties

redox-active moiety	A photoinducibly redox-active moiety or a chemically-inducibly redox-active moiety
electron donor	In the context of the present invention, the term "electron donor" refers to a component of a photoinducibly or chemically-inducibly redox-active moiety. An electron donor is a molecule that can transfer an electron to an electron acceptor, directly or under the influence of certain external conditions. One such external condition is light absorption by the electron donor or acceptor of a photoinducibly redox-active moiety. Upon irradiation with light of a specific or any given wavelength, the electron donor "D" gives up an electron to the/an electron acceptor "A" and a charge-separated state D^+A^- is formed, at least temporarily, comprising the oxidized donor and the reduced acceptor. Another such external condition may be the oxidation or reduction of the electron donor or acceptor of the chemically-inducibly redox-active moiety by an external oxidizing or reducing agent, so for example the transfer of an electron to the electron donor by a reducing agent, or the release of an electron by the electron acceptor to an oxidizing agent. These oxidizing or reducing agents may be external redox-active substances, i.e. they are not covalently joined with but in contact with the redox-active moiety, the nucleic acid oligomer, or the conductive surface, for example via the solution added to the modified conductive surface, or they

are covalently joined with the nucleic acid oligomer, the oxidizing or reducing agent being covalently attached to the nucleic acid oligomer at a site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the redox-active moiety, preferably at the oligonucleotide end opposite the modification with the redox-active moiety, near the conductive surface. The ability to act as an electron donor or acceptor is relative, i.e. a molecule that acts as an electron donor toward another molecule, directly or under the influence of certain external conditions, may also act as an electron acceptor toward that molecule under differing experimental conditions, or toward a third molecule under the same or differing experimental conditions.

electron acceptor

In the context of the present invention, the term "electron acceptor" refers to a component of a photoinducibly or chemically-inducibly redox-active moiety. An electron acceptor is a molecule that can take up an electron from an electron donor, directly or under the influence of certain external conditions. One such external condition is light absorption by the electron donor or acceptor of a photoinducibly redox-active moiety. Upon irradiation with light of a specific or any given wavelength, the electron donor "D" gives up an electron to the/an electron acceptor "A" and a charge-separated state D^+A^- is formed, at least temporarily, comprising the oxidized donor and reduced acceptor. Another such external condition may be the oxidation or reduction of the electron donor or acceptor of the chemically-inducibly redox-active moiety by an external oxidizing or reducing agent, so for example the transfer of an electron to the electron donor by a reducing agent, or the release of an electron by the

electron acceptor to an oxidizing agent. These oxidizing or reducing agents may be external redox-active substances, i.e. they are not covalently joined with but in contact with the redox-active moiety, the nucleic acid oligomer, or the conductive surface, for example via the solution added to the modified conductive surface, or they are covalently joined with the nucleic acid oligomer, the oxidizing or reducing agent being covalently attached to the nucleic acid oligomer at a site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the redox-active moiety, preferably at the oligonucleotide end opposite the modification with the redox-active moiety, near the conductive surface. The ability to act as an electron acceptor or donor is relative, i.e. a molecule that acts as an electron acceptor toward another molecule, directly or under the influence of certain external conditions, may also act as an electron donor toward that molecule under differing experimental conditions, or toward a third molecule under the same or differing experimental conditions.

electron-donor
molecule

Equivalent to an electron donor.

electron-acceptor
molecule

Equivalent to an electron acceptor.

oxidizing agent

A chemical compound (chemical substance) that oxidizes another chemical compound (chemical substance, electron donor, electron acceptor) by taking up electrons from this other chemical compound (chemical substance, electron donor, electron acceptor). An oxidizing agent behaves analogously to an electron acceptor, but is used in the context of the present invention to denote an external electron acceptor not directly belonging to the

photoinducibly or chemically-inducibly redox-active moiety. In this context, "not directly" means that the oxidizing agent is either a free redox-active substance that is not bound to but in contact with the nucleic acid oligomer, or that the oxidizing agent is covalently attached to the nucleic acid oligomer, but at a nucleic acid oligomer site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the (photoinducibly) redox-active moiety. In particular, the electrode may represent the oxidizing agent.

reducing agent

A chemical compound (chemical substance) that reduces another chemical compound (chemical substance, electron donor, electron acceptor) by giving up electrons to this other chemical compound (chemical substance, electron donor, electron acceptor). A reducing agent behaves analogously to an electron donor but is used in the context of the present invention to denote an external electron donor not directly belonging to the photoinducibly or chemically-inducibly redox-active moiety. In this context, "not directly" means that the reducing agent is either a free redox-active substance that is not bound to but in contact with the nucleic acid oligomer, or that the reducing agent is covalently attached to the nucleic acid oligomer, but at a nucleic acid oligomer site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the redox-active moiety. In particular, the electrode may represent the reducing agent.

photoinducible

Photoinducible means that a certain property is exhibited only upon irradiation with light of a specific or any given wavelength. For example, a photoinducibly redox-active moiety exhibits its redox activity, in other words its property of

	carrying out a charge separation within the photoinducibly redox-active moiety under certain external conditions, for example forming the state D^+A^- and giving up electrons to another suitable oxidizing agent or taking up electrons from another suitable reducing agent, only upon irradiation with light of a specific or any given wavelength. A further example is the photoinducibly reactive group, i.e. a group that becomes reactive only upon irradiation with light of a specific or any given wavelength.
redox-active	Redox-active refers to the property of a redox-active moiety of giving up electrons to a suitable oxidizing agent or taking up electrons from a suitable reducing agent under certain external conditions, or the property of a redox-active substance of giving up electrons to a suitable electron acceptor or taking up electrons from a suitable electron donor under certain external conditions.
free redox-active substance	A free oxidizing or reducing agent not covalently joined with but in contact with the redox-active moiety, the nucleic acid oligomer, or the conductive surface, for example via the solution added to the modified conductive surface, the free redox-active substance being able to be for example an uncharged molecule, any salt, or a redox-active protein or enzyme (oxido-reductase). The free redox-active substance is characterized in that it can rereduce the oxidized donor (or reoxidize the reduced acceptor) of the photoinducibly redox-active moiety, or in that the free redox-active substance can reduce the donor (or oxidize the acceptor) of the chemically-inducibly redox-active moiety.
photoinducibly redox-active moiety	Generic term for a moiety that includes one or more electron-donor molecules and one or more electron-acceptor

molecules, this (these) electron-donor molecule(s) and/or electron-acceptor molecule(s) being able to be embedded in one or more macromolecules. Electron donor(s) and electron acceptor(s) may be joined with one another via one or more covalent or ionic bonds, via hydrogen bonds, van der Waals bridges, π - π -interaction, or via coordination by means of electron-pair donation and acceptance, covalent links being able to be direct or indirect links (e.g. via a spacer, but not via a nucleic acid oligomer). In addition, electron donor(s) and/or electron acceptor(s) that are embedded in one or more macromolecule(s) may be joined with the macromolecule(s) via covalent attachment to the macromolecule(s), via encapsulation in suitable molecular cavities (binding pockets) of the macromolecule(s), via ionic bonds, hydrogen bonds, van der Waals bridges, π - π -interaction, or via coordination by means of electron-pair donation and acceptance between the macromolecule(s) and the electron-donor molecule(s) and/or electron-acceptor molecule(s). If the photoinducibly redox-active moiety is composed of multiple macromolecules, the binding of the macromolecules to one another may likewise take place covalently, ionically, via hydrogen bonds, van der Waals bridges, π - π -interaction, or via coordination by means of electron-pair donation and acceptance. In addition to comprising electron donor(s) and electron acceptor(s) or electron donor(s), electron acceptor(s), and macromolecule(s), essential features of the photoinducibly redox-active moiety are: (i) in the forms relevant to the present invention (electron donor(s) and electron acceptor(s) in their original state or in the oxidized or reduced state), the moiety is stable and does not dissociate into its components, (ii) the moiety includes no

nucleic acid, (iii) the moiety's composition comprising electron donor(s) and electron acceptor(s) or electron donor(s), electron acceptor(s), and macromolecule(s) can be recognized by a person skilled in the art, regardless of the bond between the components, since, in principle, they may also occur as single molecules, and (iv) under the same or similar external conditions, electron donor(s) and electron acceptor(s) of the photoinducibly redox-active moiety in the form of single molecules in solution act as electron donor(s) and electron acceptor(s), i.e. even in the case of free dissolved electron donor(s) and electron acceptor(s), an electron can be transferred from the dissolved electron donor(s) to the dissolved electron acceptor(s), directly or under the influence of certain external conditions, depending on the conditions that lead to an electron transfer within the photoinducibly redox-active moiety. The photoinducibly redox-active moiety may be for example any photoinducibly redox-active protein/enzyme or any photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex. Upon irradiation with light of a specific or any given wavelength, the/an electron donor gives up an electron to one of the electron acceptors and a charge-separated state D^+A^- is formed, at least temporarily, comprising an oxidized donor and a reduced acceptor. This process within the photoinducibly redox-active moiety is referred to as photoinduced charge separation. Given appropriately chosen external conditions, the photoinducibly redox-active moiety exhibits its redox activity, in other words its property of giving up electrons to a suitable oxidizing agent or taking up electrons from a suitable reducing agent, only in a charge-separated state, since the reducing

chemically-inducibly
redox-active moiety

agent transfers (or the oxidizing agent takes up) electrons only to the oxidized donor (or from the reduced acceptor) of the photoinducibly redox-active moiety, for example in the presence of a reducing agent that can reduce D^+ but not D (or in the presence of an oxidizing agent that can oxidize A^- but not A). In particular, this oxidizing or reducing agent may also be an electrode, the photoinducibly redox-active moiety being able to give up an electron to an electrode (or take up an electron therefrom) only subsequent to photoinduced charge separation, for example if the electrode is set to a potential at which A^- but not A is oxidized (or at which D^+ but not D is reduced).

Corresponds to a photoinducibly redox-active moiety in composition and manner of functioning, but in contrast to the manner in which a photoinducibly redox-active moiety functions, photoactivation is precluded as an external condition for developing the redox activity of the redox-active moiety. The redox-active moiety may be for example any redox-active protein/enzyme or any redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex. Given appropriately chosen external conditions, the redox-active moiety exhibits its redox activity, for example its property of giving up electrons to a suitable oxidizing agent, only after an electron is transferred from a reducing agent to the/an electron donor "D" that can transfer an electron to the acceptor "A" only in the reduced state " D^- ," and the oxidizing agent takes up electrons only from this reduced acceptor " A^- " of the redox-active moiety, in other words in the presence of an oxidizing agent that can oxidize A^- but not A (successive charge transfer). In particular, this oxidizing agent may also be an

photoinducibly
redox-active
protein/enzyme

electrode, for example if the electrode is set to a potential at which A^- but not A is oxidized. Conversely, given differently chosen external conditions, the redox-active moiety can exhibit its redox activity, for example its property of taking up electrons from a suitable reducing agent, only after an electron is transferred from an electron acceptor " A " to an oxidizing agent, if only the oxidized acceptor " A^+ " can take up an electron from the donor D and the reducing agent can transfer electrons only to the oxidized donor " D^+ " of the redox-active moiety, for example in the presence of a reducing agent that can reduce D^+ but not D (successive charge transfer). In particular, this reducing agent may also be an electrode, for example if the electrode is set to a potential at which D^+ but not D is reduced.

Usually consists of what is known as 'apoprotein' – the preferred macromolecule(s) of the present invention – and cofactors – electron donor(s) and electron acceptor(s) within the meaning of the present invention. Photoinduced charge separation within the photoactivatable redox-active protein/enzyme is triggered by light of a specific or any given wavelength. Thus, for example, in the photosynthetic reaction center (RC), cofactors in the form of a primary electron donor P and multiple different electron acceptors A , including quinone cofactor(s) Q , are embedded in a protein matrix, thus forming a "polymolecular" moiety (cf. Structure 1). In this case, the embedding takes place via encapsulation of the cofactors in suitable cavities (known as binding pockets) of the protein matrix comprising multiple protein sub-units. In the case of some naturally occurring RCs, both the protein sub-units and the encapsulation of the cofactors in the protein matrix are

chemically-inducibly
redox-active protein/
enzyme

realized via non-covalent bonds. Upon irradiation with light of a suitable wavelength, the primary donor gives up an electron to one of the electron acceptors and a charge-separated RC-state P^+A^- , and especially the state P^+Q^- , is formed, at least temporarily, comprising the initially neutral cofactors.

Corresponds to a photoinducibly redox-active protein/enzyme in composition and manner of functioning, but in contrast to the manner in which a photoinducibly redox-active protein/enzyme functions, photoactivation is precluded as an external condition for developing the redox activity of the redox-active moiety; the chemically-inducibly redox-active protein/enzyme usually consists of what is known as apoprotein – the preferred macromolecule(s) of the present invention – and cofactors – electron donor(s) and electron acceptor(s) within the meaning of the present invention. The redox-active protein's/enzyme's property of successive charge transfer is triggered by a free redox-active substance (substrate).

photoinducibly
redox-active, linked,
at least bimolecular
electron-
donor/electron-
acceptor complex

A compound comprising one or more electron-donor molecules D1, D2, D3, etc. and at least one or more suitable electron-acceptor molecules A1, A2, A3, etc., the electron donor(s) and electron acceptor(s) being joined with one another via one or more covalent or ionic bonds, via hydrogen bonds, van der Waals bridges, via π - π -interaction, or via coordination by means of electron-pair donation and acceptance. Covalent links in this sense may be direct or indirect links (e.g. via a spacer, but not via a nucleic acid oligomer). In addition to comprising electron donor(s) and electron acceptor(s), essential features of the photoinducibly redox-active, linked, at least bimolecular electron-

donor/electron-acceptor complex are: (i) in the forms relevant to the present invention (electron donor(s) and electron acceptor(s) in their original state or in the oxidized or reduced state), the electron-donor/electron-acceptor complex is stable and does not dissociate into its components, (ii) the moiety includes no nucleic acid, (iii) the at least bimolecular electron-donor/electron-acceptor complex's composition comprising electron donor(s) and electron acceptor(s) can be recognized by a person skilled in the art, regardless of the bond between the components, and (iv) under the same or similar external conditions, electron donor(s) and electron acceptor(s) of the photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex, also in the form of single molecules in solution, act as electron donor(s) and electron acceptor(s), i.e. even in the case of free dissolved electron donor(s) and electron acceptor(s), an electron can be transferred from the dissolved electron donor(s) to the dissolved electron acceptor(s), directly or under the influence of certain external conditions, depending on the conditions that lead to an electron transfer within the photoinducibly redox-active moiety. In its manner of functioning that is relevant to the present invention, the photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex corresponds to a photoinducibly redox-active protein/enzyme, i.e. here, too, irradiation with light of a suitable wavelength leads to photoinduced charge separation and, at least temporarily, a charge-separated state D^+A^- is formed (wherein D stands for any D1, D2, D3, etc. and A for any A1, A2, A3, etc.). In the expression "photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex,"

chemically-inducibly
redox-active, linked,
at least bimolecular
electron-
donor/electron-
acceptor complex

the term "at least bimolecular" indicates that the complex is composed of at least one electron donor and at least one electron acceptor, even if the donor is directly (or indirectly via a spacer) covalently joined with the acceptor.

Corresponds to a photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex in composition and manner of functioning, but in contrast to the manner in which a photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex functions, photoactivation is precluded as an external condition for developing the redox activity of the redox-active moiety. Given appropriately chosen external conditions, the redox-active moiety exhibits its redox activity, for example its property of giving up electrons to a suitable oxidizing agent, only after an electron is transferred from a reducing agent to the/an electron donor "D" that can transfer an electron to the acceptor "A" only in the reduced state "D⁻," and the oxidizing agent takes up electrons only from this reduced acceptor "A⁻" of the redox-active moiety, in other words in the presence of an oxidizing agent that can oxidize A⁻ but not A (successive charge transfer). In particular, this oxidizing agent may also be an electrode, for example if the electrode is set to a potential at which A⁻ but not A is oxidized. Conversely, given differently chosen external conditions, the redox-active moiety can exhibit its redox activity, for example its property of taking up electrons from a suitable reducing agent, only after an electron is transferred from an electron acceptor "A" to an oxidizing agent, if only the oxidized acceptor "A⁺" can take up an electron from the donor D and the reducing agent can transfer electrons only to the oxidized donor "D⁺" of the redox-active

moiety, for example in the presence of a reducing agent that can reduce D^+ but not D (successive charge transfer). In particular, this reducing agent may also be an electrode, for example if the electrode is set to a potential at which D^+ but not D is reduced. In the expression "redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex," the term "at least bimolecular" indicates that the complex is composed of at least one electron donor and at least one electron acceptor, even if the donor is directly (or indirectly via a spacer) covalently joined with the acceptor.

RC

Reaction center. An example of a photoinducibly redox-active protein/enzyme. The protein/enzyme is what is known as a pigment/protein complex composed of apoprotein comprising multiple protein sub-units and multiple cofactors (known as pigments in the example of RC). The first steps of light-driven charge separation in bacterial or vegetable photosynthesis take place in such pigment/protein complexes. For example, the RC of photosynthesizing bacteria of the strain *Rhodobacter sphaeroides* (cf. Structure 1) consists of three protein sub-units and eight cofactors (pigments). The cofactors, a bacteriochlorophyll dimer P , two bacteriochlorophyll monomers B_A and B_B , two bacteriopheophytin monomers H_A and H_B , and two ubiquinone-50 (UQ) molecules Q_A and Q_B , are localized in the respective protein binding pockets (i.e. the P , B_A , etc. binding pockets).

Q_A protein binding pocket

A protein binding pocket (or protein environment) in which the quinone cofactor Q_A is located. In the RC of *Rhodobacter sphaeroides*, for example, the quinone cofactor Q_A is a ubiquinone-50 (cf. Structure 1).

Q_A binding pocket

Q_A protein binding pocket

[019] Chemical Substances/Groups

ZnBChl	Zinc bacteriochlorophyll (Formula 11 with M = Zn)
Q	Generally quinone; in Example 3 and the passages referring thereto, Q is a modified anthraquinone or pyrroloquinoline quinone (PQQ).
UQ	Ubiquinone-50, RC cofactor and temporary electron acceptor, for example in the RC of the photosynthesizing bacteria of <i>Rhodobacter sphaeroides</i> or <i>Rhodopseudomonas viridis</i> .
(cyt c ₂) ²⁺	Reduced form of the cytochrome c ₂ , a freely movable hemoprotein that, in bacterial photosynthesis in <i>Rhodobacter sphaeroides</i> , reduces the oxidized primary donor P ⁺ to P; an example of a redox-active substance.
PQQ	Pyrroloquinoline quinone; corresponds to 4,5-dihydro-4,5-dioxo-1H-pyrrolo-[2,3-f]-quinoline-2,7,9-tricarboxylic acid
EDTA	ethylenediamine tetraacetate (sodium salt)
sulfo-NHS	N-hydroxysulfosuccinimide
EDC	(3-dimethylaminopropyl)-carbodiimide
HEPES	N-[2-hydroxyethyl]piperazine-N'-[2-ethanesulfonic acid]
Tris	trishydroxymethylamino methane
alkyl	The term "alkyl" refers to a saturated hydrocarbon group that is straight-chain or branched (e.g. ethyl, 2,5-dimethylhexyl, or isopropyl, etc.). When "alkyl" is used to indicate a linker or spacer, the term refers to a group having two available valences for covalent linkage (e.g. -CH ₂ CH ₂ -, -CH ₂ C(CH ₃) ₂ CH ₂ CH ₂ C(CH ₃) ₂ CH ₂ - or -CH ₂ CH ₂ CH ₂ -, etc.). Alkyl groups preferred as substituents or side chains R are those having a chain length of 1 – 30 (the longest continuous chain of atoms

	covalently bound to one another). Alkyl groups preferred as linkers or spacers are those having a chain length of 1 – 20, especially a chain length of 1 – 14, the chain length here representing the shortest continuous link between the structures joined via the linker or spacer, in other words between the two molecules or between a surface atom, surface molecule, or surface molecule group and another molecule.
alkenyl	Alkyl groups in which one or more of the C-C single bonds are replaced by C=C double bonds.
alkynyl	Alkyl or alkenyl groups in which one or more of the C-C single or C=C double bonds are replaced by C≡C triple bonds.
heteroalkyl	Alkyl groups in which one or more of the C-H bonds or C-C single bonds are replaced by C-N, C=N, C-P, C=P, C-O, C=O, C-S, or C=S bonds.
heteroalkenyl	Alkenyl groups in which one or more C-H bonds, C-C single or C=C double bonds are replaced by C-N, C=N, C-P, C=P, C-O, C=O, C-S, or C=S bonds.
heteroalkynyl	Alkynyl groups in which one or more of the C-H bonds, C-C single, C=C double or C≡C triple bonds are replaced by C-N, C=N, C-P, C=P, C-O, C=O, C-S, or C=S bonds.
linker	A molecular link between two molecules or between a surface atom, surface molecule, or surface molecule group and another molecule. Linkers can usually be purchased in the form of alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, or heteroalkynyl chains, the chain being derivatized in two places with (identical or different) reactive groups. These groups form a covalent chemical bond in simple/known chemical reactions with the appropriate reaction partners. The reactive groups may also be photoactivatable, i.e. the reactive groups are activated only by light of a specific or any given wavelength.

spacer	<p>Preferred linkers are those having a chain length of 1 – 20, especially a chain length of 1 – 14, the chain length here representing the shortest continuous link between the structures to be joined, in other words between the two molecules or between a surface atom, surface molecule, or surface molecule group and another molecule.</p> <p>A linker that is covalently attached via the reactive groups to one or both of the structures to be joined (see linker). Preferred spacers are those having a chain length of 1 – 20, especially a chain length of 1 – 14, the chain length representing the shortest continuous link between the structures to be joined.</p>
(n x HS-spacer)-oligo	<p>A nucleic acid oligomer to which n thiol functions are each attached via a spacer, each spacer being able to have a different chain length (the shortest continuous link between the thiol function and the nucleic acid oligomer), especially any chain length between 1 and 14. These spacers, in turn, may be bound to various reactive groups that are naturally present on the nucleic acid oligomer or that have been affixed thereto by modification, and "n" is any integer, especially a number between 1 and 20.</p>
(n x R-S-S-spacer)-oligo	<p>A nucleic acid oligomer to which n disulfide functions are each attached via a spacer, the disulfide function being saturated by any residue R. Each spacer for attaching the disulfide function to the nucleic acid oligomer may have a different chain length (shortest continuous link between the disulfide function and the nucleic acid oligomer), especially any chain length between 1 and 14. These spacers, in turn, may be bound to various reactive groups that are naturally present on the nucleic acid oligomer or that have been affixed thereto by modification. The variable n is any integer, especially a number</p>

oligo-spacer-S-S-spacer-oligo	between 1 and 20. Two identical or different nucleic acid oligomers that are joined with each other via a disulfide bridge, the disulfide bridge being attached to the nucleic acid oligomers via any two spacers, the two spacers being able to have differing chain lengths (the shortest continuous link between the disulfide bridge and the respective nucleic acid oligomer), especially any chain length between 1 and 14, and these spacers, in turn, being able to be bound to various reactive groups that are naturally present on the nucleic acid oligomer or that have been affixed thereto by modification.
-------------------------------	---

[020] Modified Surfaces/Electrodes

mica	Muscovite lamina, a support material for the application of thin films.
<i>Au-S-(CH₂)₂-ss-oligo-spacer-UQ(RC)</i>	Gold film on mica having a covalently applied monolayer of derivatized 12-bp single-strand DNA oligonucleotide (sequence: TAGTCGGAAGCA). Here, the oligonucleotide's terminal phosphate group at the 3'-end is esterified with (HO-(CH ₂) ₂ -S) ₂ to form P-O-(CH ₂) ₂ -S-S-(CH ₂) ₂ -OH, the S-S bond being homolytically cleaved and producing one Au-S-R bond each. The terminal thymine base at the 5'-end of the oligonucleotide is modified at the C-5 carbon with -CH=CH-CO-NH-CH ₂ -CH ₂ -NH ₂ , this residue, in turn, being joined via its free amino group with the carboxylic-acid group of the modified ubiquinone-50 by amidation. Thereafter, the UQ is reconstituted with the remaining RC.
<i>Au-S-(CH₂)₂-ds-oligo-spacer-UQ(RC)</i>	<i>Au-S-(CH₂)₂-ss-oligo-spacer-UQ(RC)</i> hybridized with the oligonucleotide that is complementary to the ss-oligo (sequence: TAGTCGGAAGCA).
<i>Au-S-(CH₂)₂-ss-oligo-</i>	Identical to <i>Au-S-(CH₂)₂-ss-oligo-spacer-UQ(RC)</i> with the

<i>spacer-Q-ZnBChl</i>	exception that, instead of the RC attached via UQ, Q-ZnBChl is attached as the photoinducibly redox-active moiety.
<i>Au-S-(CH₂)₂-ds-oligo-spacer-Q-ZnBChl</i>	<i>Au-S-(CH₂)₂-ss-oligo-spacer-Q-ZnBChl</i> hybridized with the oligonucleotide that is complementary to the ss-oligo (sequence: TAGTCGGAAGCA).

[021]

Electrochemistry

<i>E</i>	The electrode potential on the working electrode.
<i>E_{ox}</i>	Potential at maximum current of the oxidation of a reversible electrooxidation or electroreduction.
<i>i</i>	current density (current per cm ² of electrode surface)
cyclic voltammetry	Recording a current-voltage curve. Here, the potential of a stationary working electrode is changed linearly as a function of time, starting at a potential at which no electrooxidation or electroreduction occurs, up to a potential at which a species that is dissolved or adsorbed on the electrode is oxidized or reduced (i.e. a current flows). After running through the oxidation or reduction operation, which produces in the current-voltage curve an initially increasing current and, after reaching a maximum, a gradually decreasing current, the direction of the potential feed is reversed. The behavior of the products of electrooxidation or electroreduction is then recorded in a reverse run.
amperometry	Recording a current-time curve. Here, the potential of a stationary working electrode is set, for example by a potential jump, to a potential at which the electrooxidation or electroreduction of a dissolved or adsorbed species occurs, and the flowing current is recorded as a function of time.

[022]

The present invention is directed to a nucleic acid oligomer that is modified by chemically binding a redox-active moiety. The redox-active moiety is either a photoinducibly redox-active moiety or a chemically-inducibly redox-active moiety. Subsequent to photoinduced release of an electron to an external oxidizing agent, for example an electrode, or taking up an electron

from an external reducing agent, for example an electrode, the photoinducibly redox-active moiety may be rereduced or reoxidized by a free redox-active substance, in other words it may be restored to its original state. After giving up an electron to an external oxidizing agent, the chemically induced redox-active moiety may be reduced by an external reducing agent, for example an electrode, or after taking up an electron from an external reducing agent, be oxidized by an external oxidizing agent, for example an electrode.

[023] In the context of the present invention, a compound comprising at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine (e.g. cytosine, thymine, or uracil) or purine bases (e.g. adenine or guanine), preferably a DNA, RNA, or PNA fragment, is used as the nucleic acid oligomer. In the present invention, the term "nucleic acid" refers to any backbone of the covalently-joined pyrimidine or purine bases, such as the sugar-phosphate backbone of DNA, cDNA, or RNA, a peptide backbone of PNA, or analogous backbone structures such as a thiophosphate, a dithiophosphate, or a phosphoramidate backbone. An essential feature of a nucleic acid within the meaning of the present invention is that it can sequence-specifically bind naturally occurring cDNA or RNA. The terms "(probe) oligonucleotide," "nucleic acid," and "oligomer" are used as alternatives to the term "nucleic acid oligomer."

[024] In the context of the present invention, the term "electron acceptor" or "electron-acceptor molecule" and the term "electron donor" or "electron-donor molecule" refer to a component of a redox-active moiety.

[025] In the context of the present invention, a "redox-active moiety" is understood to be any moiety that includes one or more electron-donor molecules and one or more electron-acceptor molecules. The electron-donor molecule(s) or molecular moiety (moieties) and electron-acceptor molecule(s) or molecular moiety (moieties) of this redox-active moiety may be joined with one another via one or more covalent or ionic bonds, via hydrogen bonds, van der Waals bridges, via π - π -interaction, or via coordination by means of electron-pair donation and acceptance, the covalent bonds being able to be direct or indirect bonds (e.g. via a spacer, but not via a nucleic acid oligomer). Furthermore, the electron-donor molecule(s) and/or electron-acceptor

molecule(s) may be integrated in one or more macromolecule(s), this integration occurring via encapsulation in suitable molecular cavities (binding pockets) of the macromolecule(s), via hydrogen bonds, van der Waals bridges, π - π -interaction, or via coordination by means of electron-pair donation and acceptance between the macromolecule(s) and the electron-donor molecule(s) and/or the electron-acceptor molecule(s). Thus, in this case, the macromolecule(s) and the electron-donor molecule(s) and electron-acceptor molecule(s) form the redox-active moiety. If multiple macromolecules are components of the redox-active moiety, the binding of the macromolecules to one another may likewise take place covalently, ionically, via hydrogen bonds, van der Waals bridges, π - π -interaction, or via coordination by means of electron-pair donation and acceptance.

[026] According to the present invention, the aforementioned donor and acceptor molecules form a redox-active moiety, i.e. they are bound to one another directly or via further molecular moieties. The sole restriction on the molecules or molecular moieties joining the components of the redox-active moiety, according to the present invention, is the exclusion of nucleic acid oligomers. According to the present invention, the redox-active moiety is bound to the probe oligonucleotide as a complete moiety, multiple chemical bonds being able, of course, to be formed between the oligonucleotide and the redox-active moiety. The exclusion of nucleic acid oligomers as the molecules or molecular moieties joining the components of the redox-active moiety is intended to demonstrate clearly that it is not individual portions of the redox-active moiety that are attached at various sites of the probe oligonucleotide. Thus, the probe oligonucleotide explicitly does not represent the link between the electron-donor molecule(s) or molecular moiety (moieties) and the electron-acceptor molecule(s) or molecular moiety (moieties) of the redox-active moiety.

[027] The redox-active moiety is either a photoinducibly redox-active moiety or a chemically-inducibly redox-active moiety.

[028] In the context of the present invention, "photoinducible" means that the redox activity of the photoinducibly redox-active moiety, in other words its property of giving up electrons to a suitable oxidizing agent or taking up

electrons from a suitable reducing agent under certain external conditions, is exhibited only upon irradiation with light of a specific or any given wavelength. Upon irradiation with light of a specific or any given wavelength, the electron donor "D" gives up an electron to one of the electron acceptors "A" and a charge-separated state D^+A^- is formed, at least temporarily, comprising the oxidized donor and reduced acceptor. This process within the photoinducibly redox-active moiety is referred to as photoinduced charge separation. Given appropriately chosen external conditions, the photoinducibly redox-active moiety exhibits its redox activity only in a charge-separated state, since the reducing agent can transfer (or the oxidizing agent can take up) electrons only to the oxidized donor (or from the reduced acceptor) of the photoinducibly redox-active moiety, for example in the presence of an oxidizing agent that can oxidize A^- but not A (or in the presence of a reducing agent that can reduce D^+ but not D).

[029] In particular, the aforementioned oxidizing or reducing agent may be an electrode, the photoinducibly redox-active moiety being able to give up an electron to the electrode (or take up an electron therefrom) only subsequent to photoinduced charge separation, for example if the electrode is set to a potential at which A^- but not A is oxidized (or D^+ but not D is reduced). In addition, the oxidizing or reducing agent may be a free redox-active substance, the photoinducibly redox-active moiety being able to give up an electron to (or take up an electron from) the free redox-active substance only subsequent to photoinduced charge separation, for example if the free redox-active substance oxidizes A^- but not A (or reduces D^+ but not D).

[030] In the context of the present invention, "chemically inducible" means that the redox activity of the chemically-inducibly redox-active moiety, in other words its property of giving up electrons to a suitable oxidizing agent (or taking up electrons from a suitable reducing agent) under certain external conditions, is exhibited only subsequent to reduction (or oxidation) by an external reducing agent (or oxidizing agent). The chemically-inducibly redox-active moiety corresponds to a photoinducibly redox-active moiety in composition and manner of functioning, but in contrast to the manner in which a photoinducibly redox-active moiety functions, photoactivation is precluded as an external

condition for developing the redox activity of the redox-active moiety. Given appropriately chosen external conditions, the chemically-inducibly redox-active moiety exhibits its redox activity, for example its property of giving up electrons to a suitable oxidizing agent, only after an electron is transferred from a reducing agent to the/an electron donor "D." Only in a reduced state "D⁻" can the electron donor transfer an electron to the acceptor "A," and the oxidizing agent can take up electrons only from this reduced acceptor "A⁻" of the redox-active moiety, for example in the presence of an oxidizing agent that can oxidize A⁻ but not A (successive charge transfer). In particular, said oxidizing agent may also be an electrode, for example if the electrode is set to a potential at which A⁻ but not A is oxidized. Conversely, given differently chosen external conditions, the chemically-inducibly redox-active moiety can exhibit its redox activity, for example its property of taking up electrons from a suitable reducing agent, only after an electron is transferred from an electron acceptor "A" to an oxidizing agent. Only in an oxidized state "A⁺" can an electron acceptor take up an electron from the donor D, and the reducing agent can transfer electrons only to the oxidized donor "D⁺" of the redox-active moiety, for example in the presence of a reducing agent that can reduce D⁺ but not D (successive charge transfer). In particular, said reducing agent may also be an electrode, for example if the electrode is set to a potential at which D⁺ but not D is reduced.

[031] In addition to comprising electron donor(s) and electron acceptor(s), or electron donor(s), electron acceptor(s), and macromolecule(s), essential features of the photoinducibly or chemically-inducibly redox-active moiety are: (i) in the forms relevant to the present invention (electron donor(s) and electron acceptor(s) in their original state or in an oxidized or reduced state), the moiety is stable and does not dissociate into its components, (ii) the moiety includes no nucleic acid, (iii) the moiety's composition comprising electron donor(s) and electron acceptor(s) or electron donor(s), electron acceptor(s), and macromolecule(s) can be recognized by a person skilled in the art, regardless of the bond between the components, since, in principle, electron donor(s) and acceptor(s) may also occur as single molecules, and (iv) electron donor(s) and electron acceptor(s) of the redox-active moiety, under

the same or similar external conditions to those of their form that is relevant to the present invention, as components of the redox-active moiety, also act as electron donor(s) and electron acceptor(s) in the form of single molecules in solution, i.e. even in the case of free dissolved electron donor(s) and electron acceptor(s), an electron can be transferred from the dissolved electron donor(s) to the dissolved electron acceptor(s), directly or under the influence of certain external conditions, depending on the conditions that lead to an electron transfer within the redox-active moiety. As in the case of the electron donor(s) and electron acceptor(s) of the photoinducibly redox-active moiety, such an external condition for the free, dissolved electron donor(s) and electron acceptor(s) may be light absorption by the free, dissolved electron donor(s) and acceptor(s), the (an) electron donor "D" giving up an electron to the (an) electron acceptor "A" and a charge-separated state D^+A^- being formed, at least temporarily, comprising a free, dissolved oxidized donor and a free, dissolved reduced acceptor. Another such external condition may be – as with the electron donor(s) and electron acceptor(s) of the chemically-inducibly redox-active moiety – the transfer of an electron to the free, dissolved electron donor by a reducing agent or the release of an electron to an oxidizing agent by the free, dissolved electron acceptor.

[032]

The photoinducibly redox-active moiety may be for example any photoinducibly redox-active protein/enzyme or any photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex. In the expression "photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex," the term "at least bimolecular" means that the complex is composed of at least one electron donor and at least one electron acceptor, even if this donor and this acceptor are directly (or indirectly via a spacer) covalently joined. Upon irradiation with light of a specific or any given wavelength, the/an electron donor gives up an electron to one of the electron acceptors and a charge-separated state D^+A^- is formed, at least temporarily, comprising an oxidized donor and a reduced acceptor. This process within the photoinducibly redox-active moiety is referred to as photoinduced charge separation. Given appropriately chosen external conditions, the photoinducibly redox-active moiety exhibits its redox activity, in other words its property of giving up electrons to a suitable oxidizing agent or

taking up electrons from a suitable reducing agent, only in a charge-separated state, since the reducing agent transfers (or the oxidizing agent takes up) electrons only to the oxidized donor (or from the reduced acceptor) of the photoinducibly redox-active moiety, for example in the presence of a reducing agent that can reduce D^+ but not D (or in the presence of an oxidizing agent that can oxidize A^- but not A). In particular, this oxidizing or reducing agent may also be an electrode, the photoinducibly redox-active moiety being able to give up an electron to an electrode (or take up an electron therefrom) only subsequent to photoinduced charge separation, for example if the electrode is set to a potential at which A^- but not A is oxidized (or D^+ but not D is reduced).

[033] The chemically-inducibly redox-active moiety may be for example any chemically-inducibly redox-active protein/enzyme or any chemically-inducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex. In the expression "chemically-inducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex," the term "at least bimolecular" means that the complex is composed of at least one electron donor and at least one electron acceptor, even if this donor and this acceptor are directly (or indirectly via a spacer) covalently joined. Given appropriately chosen external conditions, the chemically-inducibly redox-active moiety exhibits its redox activity, in other words its property of giving up electrons to a suitable oxidizing agent (or taking up electrons from a suitable reducing agent) under certain external conditions, only subsequent to reduction (or subsequent to oxidation) by an external reducing agent (or oxidizing agent). Only after an electron is transferred from a reducing agent to the/an electron donor " D " can the then-reduced donor " D^- " transfer an electron to the acceptor " A ," and the oxidizing agent can take up electrons only from this reduced acceptor " A^- " of the redox-active moiety. In particular, the aforementioned oxidizing agent may also be an electrode, for example if the electrode is set to a potential at which A^- but not A is oxidized. Conversely, given differently chosen external conditions, only after an electron is transferred from the acceptor " A " to an external oxidizing agent can the electron acceptor of the chemically-inducibly redox-active moiety, in its then-oxidized state " A^+ ," take up an electron from the donor D and the reducing agent can transfer electrons only to the oxidized

donor "D⁺" of the redox-active moiety. In particular, the aforementioned reducing agent may also be an electrode, for example if the electrode is set to a potential at which D⁺ but not D is reduced.

[034] In the context of the present invention, the term "oxidizing agent" refers to a chemical compound (chemical substance) that, by taking up electrons from another chemical compound (chemical substance, electron donor, electron acceptor), oxidizes this other chemical compound (chemical substance, electron donor, electron acceptor). The oxidizing agent behaves analogously to an electron acceptor but is used in the context of the present invention to refer to an external electron acceptor not directly belonging to the redox-active moiety. In this context, "not directly" means that the oxidizing agent is either a free redox-active substance that is not bound to but in contact with the nucleic acid oligomer, or that the oxidizing agent is covalently attached to the nucleic acid oligomer, but at a nucleic acid oligomer site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the redox-active moiety. In particular, the electrode may represent the oxidizing agent.

[035] Within the context of the present invention, the term "reducing agent" refers to a chemical compound (chemical substance) that, by giving up electrons to another chemical compound (chemical substance, electron donor, electron acceptor), reduces this other chemical compound (chemical substance, electron donor, electron acceptor). The reducing agent behaves analogously to an electron donor but is used in the context of the present invention to refer to an external electron donor not directly belonging to the redox-active moiety. In this context, "not directly" means that the reducing agent is either a free redox-active substance that is not bound to but in contact with the nucleic acid oligomer, or that the reducing agent is covalently attached to the nucleic acid oligomer, but at a nucleic acid oligomer site that is situated at least two covalently-joined nucleotides or at least two covalently-joined pyrimidine or purine bases away from the covalent attachment site of the redox-active moiety. In particular, the electrode may represent the reducing agent.

[036] In the context of the present invention, the term "free redox-active substance" refers to a free oxidizing or reducing agent not covalently joined with but

in contact with the redox-active moiety, the nucleic acid oligomer, or the conductive surface, for example via the solution applied to the modified conductive surface, the free redox-active substance being able to be for example an uncharged molecule, any salt, or a redox-active protein or enzyme (oxido-reductase). The free redox-active substance is characterized in that it can rereduce the oxidized donor (or reoxidize the reduced acceptor) of the photoinducibly redox-active moiety, or in that the free redox-active substance can reduce the donor (or oxidize the acceptor) of the chemically-inducibly redox-active moiety. Furthermore, the free redox-active substance is characterized in that it is oxidizable and reducible at a potential φ , where φ satisfies the condition $2.0 \text{ V} \geq \varphi \geq -2.0 \text{ V}$. The potential refers here to the free redox-active molecule in a suitable solvent, measured against a normal hydrogen electrode. In the context of the present invention, the potential range $1.7 \text{ V} \geq \varphi \geq -1.7 \text{ V}$ is preferred, the range $1.4 \text{ V} \geq \varphi \geq -1.2 \text{ V}$ being particularly preferred and the range $0.9 \text{ V} \geq \varphi \geq -0.7 \text{ V}$, at which the redox-active substances of the application examples are oxidized (or reduced), being most particularly preferred.

[037] The modified nucleic acid oligomer is directly or indirectly (via a spacer) bound to a conductive surface. The term "conductive surface" is understood to mean any electrically conductive surface of any thickness, especially metallic surfaces, surfaces comprising metal alloys, or doped or non-doped semiconductor surfaces, all semiconductors being able to be used in the form of pure substances or in the form of mixtures. In the context of the present invention, the conductive surface may be present alone or applied to any support material such as glass. In the context of the present invention, the term "electrode" is used as an alternative to "conductive surface."

[038] The term "modified conductive surface" is understood to mean a conductive surface that is modified by attaching a nucleic acid oligomer modified with a redox-active moiety.

[039] According to a further aspect, the present invention is directed to a method that allows the electrochemical detection of molecular structures, in particular the electrochemical detection of DNA/RNA/PNA fragments in a probe solution by sequence-specific nucleic acid oligomer hybridization. The detection of the hybridization events via electrical signals is a simple and

economical method and, in a battery-operated variation, allows on-site application.

[040] The present invention further provides a photoaddressable read-out method for detecting molecular structures, inter alia for detecting hybridization events on an oligomer chip, for example via electrical signals. According to the present invention, a "photoaddressable (oligomer chip) read-out method" is understood to be a method in which the detection of molecular structures is limited to a specific test site or group of test sites within the entire system (of the complete oligomer chip) by light of a specific or any given wavelength being spatially focused on (limited to) this test site (group) in order to induce the redox activity of the photoinducibly redox-active moiety.

[041] **Binding a Redox-Active Moiety to a Nucleic Acid Oligomer**

[042] A prerequisite for the method according to the present invention is the binding of a photoinducibly redox-active moiety or a chemically-inducibly redox-active moiety to a nucleic acid oligomer.

[043] The following are some examples of a photoinducibly redox-active moiety:


[044] (i) The photosynthetic bacterial reaction center (RC), such as the RC of *Rhodobacter sphaeroides* having the schematic Structure 1; the RC of other photosynthetic bacteria, such as the reaction center of *Rhodopseudomonas viridis* or *Rhodobacter capsulatus*; or a reaction center of photosynthesizing plants, such as photosystem 1 or photosystem 2, as examples of a photoinducibly redox-active protein/enzyme.

[045] (ii) Cyclophanes, or bridged porphyrin quinone systems, having the general Structure 2, as an example of a photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex. The two spacer-bridged covalent links (" - spacer - " in Structure 2) between the electron acceptor (1,4-benzoquinone in Structure 2) and the electron donor (metalloporphyrin in Structure 2) may be affixed to any site on the electron donor and/or electron acceptor. In addition to the electron acceptors shown in Structure 2, flavins having the general Formula 1, nicotinamides having the general Formula 2 or other quinones, e.g. those having the general


Formulas 3 – 8, or organic or inorganic electron acceptors and, moreover, in addition to the (metallo)porphyrins having the general Formula 9, other electron donors such as (metallo)chlorophylls having the general Formula 10 or (metallo)bacteriochlorophylls having the general Formula 11, or other organic or inorganic electron donors may also be used. In addition, simple covalently (spacer-)bridged electron-donor/electron-acceptor complexes such as covalent compounds of a substance according to Formula 9 and one of the substances according to one of Formulas 1 – 8, covalent compounds of a substance according to Formula 10 and one of the substances according to one of Formulas 1 – 8, or covalent compounds of a substance according to Formula 11 and one of the substances according to one of Formulas 1 – 8 may be used as photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complexes.

[046] (iii) Photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complexes in which (one of) the electron donor(s) and/or (one of) the electron acceptor(s) is a charge transfer complex or transition metal complex. Examples of transition metal complexes are $[\text{Ru}(\text{bipy})_2(\text{py})(\text{im})]^{2+}$, any other $[\text{Ru}(\text{II})(\text{L1})(\text{L2})(\text{L3})(\text{L4})(\text{L5})(\text{L6})]$ complexes, Cr(III), Fe(II), Os(II), or Co(II) complexes, wherein "bipy" stands for a bispyridyl ligand, "py" for a pyridyl ligand, "im" for an imidazole ligand, and L1 to L3 for any ligand, and also more or fewer than 6 ligands may coordinate on a transition metal.


[047] Examples of a chemically-inducibly redox-active moiety are the cytochrome-bc complex or the cytochrome- c_2 complex of photosynthesizing bacteria (a complex comprising a protein matrix and four embedded iron-porphyrin cofactors as electron donors and/or electron acceptors), as examples of a chemically-inducibly redox-active protein/enzyme or, as listed under (ii) and (iii), suitably composed cyclophanes or analogous compounds, as examples of a chemically-inducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex.




[048] *Structure 1:* Reaction center consisting of the cofactors P (primary donor, a bacteriochlorophyll dimer), B_A and B_B (bacteriochlorophyll monomers), H_A and H_B (bacteriopheophytins), Q_A and Q_B (ubiquinone-50) and the protein sub-units L, M, and H (not shown) that envelop the cofactors.




[049] *Structure 2:* A cyclophane; M = e.g. 2H, Mg, Zn, Cu, Ni, Pd, Co, Cd, Mn, Fe, Sn, Pt, etc.; R₁ to R₈, or spacers, are, independently of one another, any alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, or heteroalkynyl substituents.




Formula 1




Formula 2




Formula 3




Formula 4




Formula 5



Formula 6




Formula 7




Formula 8


[050] R_1 to R_8 are, independently of one another, H or any alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, or heteroalkynyl substituents.



Formula 9



Formula 10



Formula 11

[051] M = 2H, Mg, Zn, Cu, Ni, Pd, Co, Cd, Mn, Fe, Sn, Pt, etc.; R₁ to R₁₂ are, independently of one another, H or any alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, or heteroalkynyl substituents.

[052] In addition, according to the present invention, a distinguishing feature of the redox-active moiety is that said moiety gives up electrons to an oxidizing agent that is likewise covalently attached to the nucleic acid oligomer, or takes up electrons from another reducing agent that is likewise covalently attached to the oligonucleotide, this oxidizing or reducing agent being able to be in particular an electrically conductive surface (electrode) and the redox-active moiety being able to be electrooxidized/electroreduced by applying an external voltage to this electrode in its electrochemically accessible potential range.

[053] According to the present invention, a distinguishing feature of the redox-active substance is that it can rereduce the photoinducibly redox-active moiety (or can reoxidize it) after the latter gives up an electron to another oxidizing agent that is distinct from the redox-active substance and covalently attached to the oligonucleotide (or after the latter takes up an electron from another reducing agent that is distinct from the redox-active substance and covalently attached to the oligonucleotide), or that the free redox-active substance can reduce (or oxidize) the donor (or the acceptor) of the chemically-inducibly redox-active moiety. According to the present invention, any redox-active

substance may be used for this as long as it is oxidizable and reducible at a potential ϕ that satisfies the condition $2.0 \text{ V} \geq \phi \geq -2.0 \text{ V}$ and as long as the potential is suitable for rereducing (or reoxidizing) said photoinducibly redox-active moiety after the latter gives up an electron to another oxidizing agent that is likewise covalently attached to the nucleic acid oligomer (or takes up an electron from another reducing agent that is likewise covalently attached to the nucleic acid oligomer), or for reducing or oxidizing said chemically-inducibly redox-active moiety. The potential refers here to the free, unmodified redox-active substance in a suitable solvent, measured against a normal hydrogen electrode. In the context of the present invention, the potential range $1.7 \text{ V} \geq \phi \geq -1.7 \text{ V}$ is preferred, the range $1.4 \text{ V} \geq \phi \geq -1.2 \text{ V}$ being particularly preferred and the range $0.9 \text{ V} \geq \phi \geq -0.7 \text{ V}$, in which the redox-active substances of the application example are oxidized (and rereduced), being most particularly preferred. Suitable possibilities are, in addition to the usual organic and inorganic redox-active molecules such as hexacyanoferrates, ferrocenes, cobaltocenes, and quinones, most importantly ascorbic acid (or the Na^+ salt thereof), $[\text{Ru}(\text{NH}_3)_6]^{2+}$, or cytochrome c_2 ($\text{cyt } c_2$) $^{2+}$, a freely movable iron-containing protein that reduces the oxidized primary donor P^+ in the RC of *Rhodobacter sphaeroides* to P and, in doing so, is itself oxidized to ($\text{cyt } c_2$) $^{3+}$.

[054] In a preferred embodiment of the present invention, the photoinducibly or chemically-inducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex is embedded in one or more macromolecules in such a way that the macromolecule acts as an electrically insulating, enveloping end of the redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex by preventing direct electrooxidation/electroreduction of the redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex at the electrode, for example in the case of direct contact between the electrode and the redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex, but allowing indirect electrooxidation/electroreduction of the electron-donor/electron-acceptor complex mediated by a double-stranded nucleic acid oligomer. Such a macromolecule can be for example a tailored cyclodextrin

that, due to its being shaped as a cut-off cone that is hollow inside, coats a cyclophane or similar electron-donor/electron-acceptor complex.

[055] According to the present invention, a redox-active moiety is covalently bound to a nucleic acid oligomer by the reaction of the nucleic acid oligomer with the redox-active moiety or portions thereof (see also the section "Manner of Executing the Invention"). This bond can be achieved in four different ways:

[056] a) A free phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine group of the oligonucleotide backbone, especially a group at one of the two ends of the oligonucleotide backbone, is used as the reactive group for forming a bond at the nucleic acid oligomer. The free, terminal phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine groups exhibit increased reactivity and thus easily undergo typical reactions such as amidation with (primary or secondary) amino groups or with acid groups; esterification with (primary, secondary, or tertiary) alcohols or with acid groups; thioester formation with (primary, secondary, or tertiary) thioalcohols or with acid groups, or condensation of amine and aldehyde with subsequent reduction of the resultant $\text{CH}=\text{N}$ bond to a $\text{CH}_2\text{-NH}$ bond. The coupling group (acid, amine, alcohol, thioalcohol, or aldehyde function) required to covalently attach the redox-active moiety is either naturally present on the redox-active moiety or is obtained by chemically modifying the redox-active moiety. The attachment of the redox-active moiety may take place completely or in portions of the moiety with subsequent completion of the redox-active moiety (see below).

[057] b) The nucleic acid oligomer is modified with a reactive group at the oligonucleotide backbone or at a base via a covalently-attached molecular moiety (spacer) of any composition and chain length (longest continuous chain of atoms bound to one another), especially a chain length of 1 to 14. The modification preferably takes places at one of the ends of the oligonucleotide backbone or at a terminal base. An alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, or heteroalkynyl substituent, for example, may be used as the spacer. Possible simple reactions for forming the covalent bond between the redox-active moiety and the nucleic acid oligomer thus modified are, as described under a), amidation from an acid and amino group, esterification from an acid and alcohol group, thioester formation from an acid and

thioalcohol group, or condensation of aldehyde and amine with subsequent reduction of the resultant $\text{CH}=\text{N}$ bond to a $\text{CH}_2\text{-NH}$ bond. The attachment of the redox-active moiety may take place completely or in portions of the redox-active moiety with subsequent completion of the moiety (see below).

[058] c) If the nucleic acid oligomer is synthesized, a terminal base will be replaced by the redox-active moiety. This attachment of the redox-active moiety may take place completely or in portions of the moiety with subsequent completion of the redox-active moiety (see below).

[059] d) If a linked (at least bimolecular) electron-donor/electron-acceptor complex is used as the redox-active moiety, the electron acceptor (or donor) will, in a first covalent modification, as described under b) or c) in this section, be bound to a terminal base, or in place of a terminal base, to the nucleic acid oligomer and thereafter, in a second covalent modification of the electron donor (or acceptor), as described under a) in this section, bound at the same end of the nucleic acid oligomer backbone to a reactive group of the backbone or to a reactive group of the acceptor (or donor). If a covalently-linked, trimolecular or greater electron-donor/electron-acceptor complex is used, then, instead of the electron acceptor (or donor), any portion of the electron-donor/electron-acceptor complex may also be used in the first covalent modification and completed in a second or further covalent modification(s).

[060] According to the present invention, the binding of the redox-active moiety to the nucleic acid oligomer may take place completely or in portions, before or after the nucleic acid oligomer is bound to the conductive surface. Thus, in the case of a redox-active protein/enzyme comprising apoprotein and cofactor(s), instead of the complete redox-active moiety, it is also possible for only the apoprotein, the apoprotein and a portion of the cofactors, or one or more cofactors to be attached and the redox-active moiety will be completed by subsequent reconstitution with the remaining missing portions. If a linked (at least bimolecular) electron-donor/electron-acceptor complex is used as the redox-active moiety, the electron acceptor (or donor) will, in a first covalent modification, as described under b) or c) in this section, be bound to a terminal base, or in place of a terminal base, to the nucleic acid oligomer and thereafter, in a second covalent modification of the electron donor (or

acceptor), as described under a) in this section, bound at the same end of the nucleic acid oligomer backbone to a reactive group of the backbone. If a covalently-linked, trimolecular or greater electron-donor/electron-acceptor complex is used, then, instead of the electron acceptor (or donor), any portion of the electron-donor/electron-acceptor complex may also be used in the first covalent modification and completed in the second covalent modification. These modifications may take place before or after the nucleic acid oligomer is bound to the conductive surface.

[061] If there are multiple different nucleic acid oligomer combinations (test sites) on a shared surface, and the redox-active moiety is to be attached to the surface after the nucleic acid oligomer is immobilized, it is advantageous to standardize the (covalent) attachment of the redox-active moiety to the nucleic acid oligomers for the entire surface by the appropriate choice of reactive group at the free nucleic acid oligomer ends of the various test sites.

[062] If redox-active proteins/enzymes are used as the redox-active moiety, the covalent attachment of the nucleic acid oligomer may take place at any reactive group that is naturally present on or affixed to the protein by modification, or – in the event that the redox-active protein/enzyme consists of apoprotein and cofactor(s) – at any reactive group that is naturally present on or affixed to a (any) cofactor by modification. In the context of the present invention, the covalent attachment at any reactive group that is naturally present on or affixed by modification to a (any) cofactor of the protein is preferred. Without wanting to be bound to mechanistic details, if there are multiple cofactors, special preference is given to the one that can give up electrons to an external oxidizing agent that is likewise covalently attached to the nucleic acid oligomer, or that can take up electrons from an external reducing agent that is likewise covalently attached to the nucleic acid oligomer (see also the section "Method of Amperometrically Detecting Nucleic Acid Oligomer Hybrids").

[063] **The Conductive Surface**

[064] According to the present invention, the term "conductive surface" is understood to mean any support having an electrically conductive surface of

any thickness, especially surfaces comprising platinum, palladium, gold, cadmium, mercury, nickel, zinc, carbon, silver, copper, iron, lead, aluminum, and manganese.

[065] In addition, any doped or non-doped semiconductor surfaces of any thickness may also be used. All semiconductors may be used in the form of pure substances or in the form of mixtures. Examples include, but are not limited to, carbon, silicon, germanium, α tin, and Cu(I) and Ag(I) halides of any crystal structure. Also suitable are all binary compounds of any composition and any structure comprising the elements of groups 14 and 16, the elements of groups 13 and 15, and the elements of groups 15 and 16. In addition, ternary compounds of any composition and any structure comprising the elements of groups 11, 13, and 16 or the elements of groups 12, 13, and 16 may be used. The designations of the groups of the periodic system refer to the IUPAC recommendation of 1985.

[066] **Binding a Nucleic Acid Oligomer to the Conductive Surface**

[067] According to the present invention, a nucleic acid oligomer is linked directly or via a linker/spacer with the surface atoms or molecules of a conductive surface of the type described above. This binding may be carried out in three different ways:

[068] a) The surface is modified in such a way that a reactive molecule group is accessible. This may take place by direct derivatization of the surface molecules, for example by wet chemical or electrochemical oxidation/reduction. Thus, for example, the surface of graphite electrodes can be provided with aldehyde or carboxylic-acid groups by wet chemical oxidation. Electrochemically, it is possible, for example by reduction in the presence of aryl-diazonium salts, to couple the appropriate (functionalized, i.e. provided with a reactive group) aryl radical, or by oxidation in the presence of $R'CO_2H$, to couple the (functionalized) R' -radical to the graphite electrode surface. An example of direct modification of semiconductor surfaces is the derivatization of silicon surfaces to reactive silanols, i.e. silicon supports having $Si-OR''$ groups on the surface, both R'' and R' representing any functionalized organic residue (e.g. alkyl, alkenyl, alkynyl, heteroalkyl,

heteroalkenyl, or heteroalkynyl substituent). Alternatively, the entire surface may be modified by covalently attaching a reactive group of a bifunctional linker, such that a monomolecular layer comprising any molecules and including a reactive group, preferably terminally, results on the surface. The term "bifunctional linker" is understood to mean any molecule of any chain length, especially chain lengths 2 – 14, having two identical (homobifunctional) or two different (heterobifunctional) reactive molecule groups.

[069] If multiple different test sites are to be formed on the surface by making use of the methodology of photolithography, then at least one of the reactive groups of the homo- or heterobifunctional linkers is a photoinducibly reactive group, i.e. a group that becomes reactive only upon irradiation with light of a specific or any given wavelength. This linker is applied in such a way that the/a photoactivatable reactive group is available after the linker is covalently attached to the surface. The nucleic acid oligomers are covalently attached to the surface thus modified, and are themselves modified with a reactive group via a spacer of any composition and chain length, especially chain lengths 1 – 14, preferably near an end of the nucleic acid oligomer. The reactive group of the oligonucleotide is one of any of the groups that react directly (or indirectly) with the modified surface to form a covalent bond. In addition, a further reactive group may be bound to the nucleic acid oligomers near their second end, this reactive group, in turn, being attached, as described above, directly or via a spacer of any composition and chain length, especially a chain length of 1 – 14. Furthermore, as an alternative to this further reactive group, the redox-active moiety (completely or portions thereof) may be attached to this second end of the nucleic acid oligomer.

[070] b) The nucleic acid oligomer that is to be applied to the conductive surface is modified with one or more reactive groups via a covalently-attached spacer of any composition and chain length, especially a chain length of 1 – 14, the reactive groups being located preferably near an end of the nucleic acid oligomer. The reactive groups are groups that can react directly with the unmodified surface. Some examples are: (i) thiol- (HS-) or disulfide- (S-S-) derivatized nucleic acid oligomers having the general formula (n x HS-spacer)-oligo, (n x R-S-S-spacer)-oligo, or oligo-spacer-S-S-spacer-oligo that react

with a gold surface to form gold-sulfur bonds or (ii) amines that attach to platinum or silicon surfaces by chemisorption or physisorption. In addition, a further reactive group may be bound to the nucleic acid oligomers near their second end, this reactive group, in turn, being attached, as described above, directly or via a spacer of any composition and chain length, especially a chain length of 1 – 14. Furthermore, as an alternative to this further reactive group, the photoinducibly redox-active moiety (completely or portions thereof) may be attached at this second end of the oligonucleotide. Particularly nucleic acid oligomers that are modified with multiple spacer-bridged thiol or disulfide bridges ((n x HS-spacer)-oligo or (n x R-S-S-spacer)-oligo) have the advantage that such nucleic acid oligomers can be applied to the conductive surface at a specific setting angle (angle between the surface normal and the helix axis of a double-stranded helical nucleic acid oligomer or between the surface normal and the axis perpendicular to the base pairs of a double-stranded non-helical nucleic acid oligomer) if the spacers attaching the thiol or disulfide functions to the nucleic acid oligomer possess an increasing or decreasing chain length as viewed from an end of the nucleic acid.

[071] c) Groups used as the reactive group on the probe nucleic acid oligomer are phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine groups of the oligonucleotide backbone, especially terminal groups. The phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine groups exhibit increased reactivity and consequently enter easily into typical reactions such as amidation with (primary or secondary) amino or acid groups, esterification with (primary, secondary, or tertiary) alcohols or acid groups, thioester formation with (primary, secondary, or tertiary) thioalcohols or acid groups, or condensation of amine and aldehyde with subsequent reduction of the resultant CH=N bond to a CH₂-NH bond. In this case, the coupling group required for covalent attachment to the phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine group is part of the surface derivatization with a (monomolecular) layer having any molecule length, as described under a) in this section, or the phosphoric-acid, sugar-C-3-hydroxy, carboxylic-acid, or amine group can react directly with the unmodified surface, as described under b) in this section. In addition, a further reactive group may be bound to

the oligonucleotides near their second end, this reactive group, in turn, being attached, as described above, directly or via a spacer of any composition and chain length, especially a chain length of 1 - 14. Furthermore, as an alternative to this further reactive group, the redox-active moiety (completely or portions thereof) may be attached to this second end of the nucleic acid oligomer.

[072] Binding the nucleic acid oligomer to the conductive surface may take place before or after the redox-active moiety is attached to the nucleic acid oligomer. In the case of a redox-active protein/enzyme comprising apoprotein and cofactor(s), instead of the complete redox-active moiety, it is also possible for only the apoprotein, the apoprotein and a portion of the cofactors, or one or more of the cofactors to be attached and the redox-active moiety will be completed by subsequent reconstitution with the remaining missing portions. If a linked (at least bimolecular) electron-donor/electron-acceptor complex is used as the redox-active moiety, the electron acceptor (or donor) may, as described under b) or c) in the section "Binding a Redox-Active Moiety to a Nucleic Acid Oligomer," be attached to a terminal base, or in place of a terminal base, to the nucleic acid oligomer, and the electron donor (or acceptor) may be attached by subsequent covalent attachment to a reactive group of the electron acceptor (or donor) or, as described under a) in the section "Binding a Redox-Active Moiety to a Nucleic Acid Oligomer," by subsequent attachment to a terminal reactive group of the nucleic acid oligomer backbone at the same end (see also the section "Manner of Executing the Invention"). Alternatively, binding the nucleic acid oligomer to the conductive surface may take place before or after the spacer having a reactive group for binding the redox-active moiety is attached. Binding the already modified nucleic acid oligomer to the conductive surface, i.e. binding to the surface after the redox-active moiety is attached to the nucleic acid oligomer or after portions of the redox-active moiety are attached, or after the spacer having a reactive group for binding the redox-active moiety is attached, likewise takes place as described under a) to c) in this section.

[073] In producing the test sites, when attaching the single-strand nucleic acid oligomers to the surface, care must be taken that sufficient distance

remains between the individual nucleic acid oligomers to provide, first, the space necessary for hybridization with the target nucleic acid oligomer and, second, the space necessary for the attachment of the redox-active moiety. Three different methods of proceeding (and combinations thereof) offer themselves for this purpose:

- [074]** 1.) Producing a modified surface by attaching a hybridized nucleic acid oligomer, in other words a surface derivatization with hybridized probe nucleic acid oligomer instead of with single-strand probe oligonucleotide. The nucleic acid oligomer strand used for hybridization is unmodified (the surface attachment is carried out as described under a) - c) in this section). Thereafter, the hybridized nucleic acid oligomer double-strand is thermally dehybridized, thus producing a single-strand-nucleic-acid-oligomer-modified surface having greater distance between the probe nucleic acid oligomers.
- [075]** 2.) Producing a modified surface by attaching a single-strand or double-strand nucleic acid oligomer, adding, during surface derivatization, a suitable monofunctional linker that, in addition to the single-strand or double-strand nucleic acid oligomer, is also bound to the surface (the surface attachment is carried out as described under a) - c) in this section). According to the present invention, the monofunctional linker has a chain length that is identical to the chain length of the spacer between the surface and the nucleic acid oligomer, or that differs by a maximum of four chain atoms. If double-strand nucleic acid oligomer is used for surface derivatization, the nucleic acid oligomer double-strand is thermally dehybridized after the double-strand nucleic acid oligomer and the linker are jointly attached to the surface. By simultaneously attaching a linker to the surface, the distance between the single-strand or double-strand nucleic acid oligomers that are likewise bound to the surface is increased. If a double-strand nucleic acid oligomer is used, this effect is amplified further by the subsequent thermal dehybridization.
- [076]** 3.) Producing a modified surface by attaching a single-strand or double-strand oligonucleotide to which the redox-active moiety is already attached, the redox-active moiety having a diameter of greater than 30 Å. If double-strand oligonucleotide is used, the oligonucleotide double-strand is thermally dehybridized after the double-strand oligonucleotide is attached to the surface.

[077] Regarding the individual steps in "Binding a Redox-Active Moiety to a Nucleic Acid Oligomer," as well as in "Binding an Oligonucleotide to the Conductive Surface," it should be noted that, in the section "Manner of Executing the Invention," the various combination possibilities of the individual steps that lead to the same end result are demonstrated in an example (Figure 2).

[078] Method of Electrochemically Detecting Nucleic Acid Oligomer Hybrids

[079] Advantageously, according to the method of electrochemically detecting nucleic acid oligomer hybrids, multiple probe nucleic acid oligomers varying in sequence, ideally all necessary combinations of the nucleic acid oligomer, are applied to an oligomer (DNA) chip to detect the sequence of any target nucleic acid oligomer or (fragmented) target DNA, or in order to seek and sequence-specifically detect mutations in the target. For this purpose, the surface atoms or molecules of a defined area (a test site) on a conductive surface are linked with DNA/RNA/PNA nucleic acid oligomers having a known but arbitrary sequence, as described above. In a most general embodiment, however, the DNA chip may also be derivatized with a single probe oligonucleotide. Preferred probe nucleic acid oligomers are nucleic acid oligomers (e.g. DNA, RNA, or PNA fragments) of base length 3 to 50, preferably of length 5 to 30, particularly preferably of length 8 to 25. According to the present invention, a redox-active moiety is or becomes bound to the probe nucleic acid oligomers, as described below.

[080] The modification of the probe nucleic acid oligomers with a redox-active moiety may take place completely or in components of the redox-active moiety, either before or after the oligonucleotide probe is bound to the conductive surface. The various combination possibilities of the individual steps (reaction sequences) are demonstrated in the section "Manner of Executing the Invention" with the aid of Figure 2 using the example of a redox-active moiety bound to an electrode via a probe oligonucleotide.

[081] Regardless of the respective reaction sequence, a surface hybrid having the general structure elec-spacer-ss-oligo-spacer-moiety develops, "moiety" representing the photoinducibly or chemically-inducibly redox-active moiety. The bridges may, of course, also be produced without spacers or with

only one spacer (elec-ss-oligo-spacer-moiety or elec-spacer-ss-oligo-moiety). In the example in Figure 2, the moiety is a photoinducibly redox-active moiety, namely the reaction center (RC) of the photosynthesizing bacteria of the strain *Rhodobacter sphaeroides*, a photoinducibly redox-active protein consisting of apoprotein and cofactors. In the example in Figures 2, 3, and 4, the RC, via its cofactor ubiquinone-50 (UQ) in what is known as the Q_A protein binding pocket of the RC, is covalently joined with the nucleic acid oligomer. The RC forms a 1:1 complex with the cofactor ubiquinone-50 in the Q_A binding pocket, the ubiquinone-50 being covalently bound to the nucleic acid oligomer in the manner described. In the example in Figures 5 and 6, the moiety is a photoinducibly redox-active, linked, at least bimolecular electron-donor/electron-acceptor complex, namely a covalently-linked zinc-bacteriochlorophyll-quinone complex that is covalently joined (via a spacer) with the nucleic acid oligomer via the quinone, the electron-acceptor molecule of the complex.

[082] The electrochemical communication between the (conductive) surface and the redox-active moiety ("moiety") bridged via a single-strand oligonucleotide having the general structure elec-spacer-ss-oligo-spacer-moiety is weak or nonexistent.

[083] In a next step, the test sites are brought into contact with the nucleic acid oligomer solution to be examined (target). This leads to hybridization only if the solution contains nucleic acid oligomer strands that are complementary to the probe nucleic acid oligomers bound to the conductive surface, or complementary in at least wide areas. Hybridization between the probe and target nucleic acid oligomers leads to increased conductivity between the surface and the redox-active moiety, since the latter is now bridged via the nucleic acid oligomer consisting of a double-strand. Figure 3 illustrates this schematically using elec-spacer-ss-oligo-spacer-UQ(RC) as an example. In Figure 4, the sequence of the electron transfer steps in elec-spacer-ds-oligo-spacer-UQ(RC) is shown in detail, while Figure 5 schematically illustrates the example elec-spacer-ss-oligo-spacer-Q-ZnBChl and Figure 6 shows in detail the sequence of the electron transfer steps in elec-spacer-ds-oligo-spacer-Q-ZnBChl.

- [084]** As a result of the hybridization of the probe nucleic acid oligomer and the nucleic acid oligomer strand that is complementary thereto (target), the electrical communication between the (conductive) surface and the photoinducibly redox-active moiety changes. Thus, a sequence-specific hybridization event can be detected by electrochemical methods such as cyclic voltammetry, amperometry, or conductivity measurements.
- [085]** In cyclic voltammetry, the potential of a stationary working electrode is changed linearly as a function of time. Starting at a potential at which no electrooxidation or electroreduction occurs, the potential is changed until the redox-active substance is oxidized or reduced (i.e., a current flows). After running through the oxidation or reduction operation, which produces in the current-voltage curve an initially increasing current, then a maximum current (peak), and finally a gradually decreasing current, the direction of the potential feed is reversed. The behavior of the products of electrooxidation or electroreduction is then recorded in a reverse run.
- [086]** An alternative electrical detection method, amperometry, is made possible by applying a suitable constant electrode potential such that the redox-active moiety may be electrooxidized (electroreduced), but the rereduction (reoxidation) of the redox-active moiety to its original state takes place, not by changing the electrode potential as in cyclic voltammetry, but rather by adding a suitable reducing agent (oxidizing agent), the "redox-active substance," to the target solution, thereby closing the current circuit of the entire system. As long as such a reducing agent (oxidizing agent) is present, or as long as the consumed reducing agent (oxidizing agent) is rereduced (reoxidized) on the counter electrode, a current flows that can be amperometrically detected and that is proportional to the number of hybridization events.
- [087]** This principle of amperometric detection will be explained in greater detail using the example of glucose oxidase to represent a photoinducibly redox-active moiety or a redox-active moiety. Glucose oxidase is a redox-active enzyme consisting of apoprotein and one cofactor (flavin adenine dinucleotide). The probe oligonucleotide having one end covalently attached to the electrode can be functionalized at the other, free end with the complete glucose oxidase enzymatic moiety, for example by covalently attaching the flavin adenine dinucleotide (FAD)

cofactor of the enzyme to the probe oligonucleotide and subsequently reconstituting it with the glucose oxidase apoprotein (GOx). The resultant surface hybrid having the general structure elec-spacer-ss-oligo-spacer-FAD(GOx) exhibits little or no conductivity between the electrode and the FAD. In the case of hybridization with the "ss-oligo"-complementary target oligonucleotide, the conductivity is significantly increased. Upon adding the glucose substrate to the target oligonucleotide solution, the FAD of the glucose oxidase (FAD(GOx)) is reduced to FADH₂ of the glucose oxidase (FADH₂(GOx)), glucose being oxidized to gluconic acid. If a suitable external potential is then applied to the electrode such that electrons from FADH₂(GOx) are given up to the electrode via the hybridized oligonucleotide, and FADH₂(GOx) is thus reoxidized to FAD(GOx) (but neither glucose nor gluconic acid can be electrooxidized or electroreduced at this potential), a current will flow in the elec-spacer-ds-oligo-spacer-FAD(GOx) system as long as FAD(GOx) is reduced by free glucose, i.e. until all of the glucose is consumed or, in the event that a potential at which gluconic acid can be reduced to glucose is applied to the counter electrode, as long as gluconic acid is reduced on the counter electrode. This current can be detected amperometrically and is proportional to the number of hybridization events.

[088] The photoinducibly or chemically-inducibly redox-active moieties of relevance to the present invention possess, however, instead of one electron donor or electron acceptor, at least one electron donor and at least one electron acceptor.

[089] In the case of a chemically-inducibly redox-active moiety, within the meaning of the present invention, at least one charge transfer step intervenes between electron donor(s) and electron acceptor(s). The free redox-active substance that reduces D (or oxidizes A) and thus initiates an electron transfer from D⁻ to A to form A⁻ (or an electron transfer from D to A⁺ to form D⁺), makes it possible to set the electrode to a potential at which A⁻ (or D⁺), but not A (or D) can be oxidized (or reduced). This has the advantage that the electrode possesses a potential at which the direct reaction of the free redox-active substance with the electrode can be significantly suppressed, and primarily electron transfers between the redox-active moiety and the electrode can be detected.

[090] If the redox-active moiety is a photoinducibly redox-active moiety, the redox activity of the moiety is triggered only by light of a specific or any given wavelength. According to the present invention, this property is used to advantage in that electrochemical detection is triggered only by radiating light onto the surface hybrid having the general structure elec-spacer-ds-oligo-spacer-moiety (surface hybrid with hybridized target) and is maintained, at most, as long as light irradiation continues. Thus, particularly in the case of amperometric detection, if a photoinducibly redox-active moiety is used, under certain external conditions, (rather long-lasting) current will flow only if light is radiated onto the surface hybrid. Such external conditions are for example the presence of a reducing agent (or oxidizing agent) suitable for reducing (or oxidizing) a photoinductively-formed oxidized donor D^+ (or reduced acceptor A^-) of the photoinducibly redox-active moiety, and applying to the electrode a potential at which a photoinductively-formed reduced acceptor A^- (or oxidized donor D^+) of the photoinducibly redox-active moiety can be oxidized (or reduced), but the non-reduced acceptor A (or the non-oxidized donor D) cannot be oxidized (or reduced). In the section "Manner of Executing the Invention," this is explained in greater detail using various examples of an elec-spacer-ss-oligo-spacer-moiety having a photoinducibly redox-active moiety. In this way, detection using a photoinducibly redox-active moiety can be spatially limited to a certain test site or group of test sites of the oligomer chip by restricting the light to this test site or group of test sites. According to the present invention, various test sites (nucleic acid oligomer combinations) of an oligomer chip can thus be applied to a shared, continuous, electrically-conductive surface. A particular test site or group of test sites can be addressed and amperometrically detected simply by applying a suitable external potential to the (entire) surface if precisely this test site or group of test sites is irradiated with light. The various test sites thus need not be applied to individual (micro-) electrodes that are electrically isolated from one another and individually controllable for applying a potential and reading out the current. Moreover, if surface hybrids having the general structure elec-spacer-ss-oligo-spacer-moiety are used with a photoinducibly redox-active moiety and amperometric detection, the read-out process for detecting individual sequence-specific hybridization events on the oligomer chip can be optimized by first reading out the test sites by

roughly scanning them with appropriately focused light and then successively increasing the resolution capacity in the grids having hybridization events, so for example, for an octamer chip having 65,536 test sites, e.g. 64 groups of 1024 test sites each are read out, then the test site groups that are shown by amperometric measurements to exhibit hybridization events can be tested e.g. in 32 groups of 32 test sites each, and thereafter, in the test site groups that again exhibit hybridization events, the test sites are assayed individually. In this way, the individual hybridization events can be quickly assigned to specific probe oligomers with little experimental outlay.

[091] BRIEF DESCRIPTION OF THE DRAWINGS

[092] The invention will be explained in greater detail below by reference to exemplary embodiments in association with the drawings, wherein:

[093] Fig. 1 Shows a schematic diagram of oligonucleotide sequencing by hybridization on a chip;

[094] Fig. 2 Shows various reaction sequences for producing the surface hybrid elec-spacer-ss-oligo-spacer-UQ(RC). The photoinducibly redox-active moiety in this surface hybrid is the reaction center (RC) of the photosynthesizing bacteria *Rhodobacter sphaeroides*. This photoinducibly redox-active protein consists of apoprotein and cofactors. The RC, via its cofactor ubiquinone-50 (UQ) in what is known as the Q_A protein binding pocket, is covalently joined via a spacer with the oligonucleotide;

[095] Fig. 3 Shows a schematic diagram of the photoinduced amperometric measurement method using the example of the surface hybrid elec-spacer-ss-oligo-spacer-UQ(RC) in Figure 2 ($h\nu$: irradiation with light; P: primary donor of the RC; UQ: ubiquinone-50 electron acceptor in the Q_A protein binding pocket of the RC; Red/Ox: reduced or oxidized form of the free redox-active substance added to the target solution, e.g. cyt c_2^{2+} , sodium ascorbate or $Fe(CN)_6^{2+}$, which can rereduce the oxidized form P^+ to its original neutral state P; E_{ox} : potential of the electrode at which UQ^- is oxidized to UQ via electron transfer to the electrode; "hv on": beginning of light irradiation; "hv off": end of light irradiation);

[096] Fig. 4 Shows a detailed schematic diagram of the surface hybrid Au-S(CH₂)₂-ds-oligo-spacer-Q-ZnBChl of Figure 3 having gold as the solid support material, mercaptoethanol as the spacer (-S-CH₂CH₂- spacer) between the electrode and the oligonucleotide, and -CH₂-CH=CH-CO-NH-CH₂-CH₂-NH- as the spacer between the electron acceptor PQQ and the oligonucleotide, as well as a diagram of the sequence of the photoinduced electron transfer steps. The apoprotein of the RC is indicated only as a shell (solid line) (cf. Structure 1). The 12-bp probe oligonucleotide of the exemplary sequence 5'-TAGTCGGAAGCA-3' in the hybridized state is shown in detail;

[097] Fig. 5 Shows a schematic diagram of the photoinduced amperometric measurement method using the example of the surface hybrid elec-spacer-ss-oligo-spacer-Q-ZnBChl ($h\nu$: irradiation with light; ZnBChl: electron-donor molecule zinc-bacteriochlorophyll; Q: the electron-acceptor molecule quinone, e.g. modified anthraquinone or PQQ; Red/Ox: reduced or oxidized form of the free redox-active substance added to the target solution, e.g. $Fe(CN)_6^{2+}$, which can rereduce the oxidized form of the electron donor $ZnBChl^+$ to its original neutral state ZnBChl; E_{ox} : electrode potential at which Q^- is oxidized to Q by giving up an electron to the electrode; "hv on": beginning of light irradiation; "hv off": end of light irradiation);

[098] Fig. 6 Shows a detailed schematic diagram of the surface hybrid Au-S(CH₂)₂-ds-oligo-spacer-Q-ZnBChl of Figure 5 having gold as the solid support material, mercaptoethanol as the spacer (-S-CH₂CH₂- spacer) between the electrode and the oligonucleotide, and -CH₂-CH=CH-CO-NH-CH₂-CH₂-NH- as the spacer between the electron acceptor PQQ and the oligonucleotide, as well as a diagram of the sequence of the photoinduced electron transfer steps. The 12-bp probe oligonucleotide of the exemplary sequence 5'-TAGTCGGAAGCA-3' in the hybridized state is shown in detail.

[099]

MANNER OF EXECUTING THE INVENTION

[100]

A formation unit of an exemplary test site with hybridized target, Au-S(CH₂)₂-ds-oligo-spacer-UQ(RC) having the general structure elec-spacer-ds-oligo-spacer-moiety, is illustrated in Figure 4. In the context of the present invention, "formation unit" is understood to mean the smallest repeating unit of a test site. In the example in Figure 4, the surface is a gold electrode. The link between the gold electrode and the probe oligonucleotide was formed with the linker (HO-(CH₂)₂-S)₂, which was esterified with the terminal phosphate group at the 3'-end to form P-O-(CH₂)₂-S-S-(CH₂)₂-OH and, following homolytic cleavage of the S-S bond at the gold surface, produced one Au-S bond each, with which 2-hydroxy-mercaptoethanol and mercaptoethanol-bridged oligonucleotide was coadsorbed on the surface. The photoinducibly redox-active moiety in the example in Figure 4 is the reaction center (RC) of the photosynthesizing bacteria *Rhodobacter sphaeroides*, a photoinducibly redox-active protein consisting of apoprotein and cofactors. In the application example, the RC, via its cofactor ubiquinone-50 (UQ) in what is known as the Q_A binding pocket of the RC, is covalently joined with the oligonucleotide, where free UQ was first provided with a reactive carboxylic-acid group (see Example 1), then covalently attached to the probe oligonucleotide via this carboxylic-acid group (amidation and dehydration of the terminal amino function of the -CH=CH-CO-NH-CH₂-CH₂-NH₂ linker attached at the C-5 position of the 5' thymine),

and finally, the remaining RC (apoprotein with all cofactors except UQ) was reconstituted to UQ.

[101] As already mentioned above, the modification of the probe oligonucleotides may take place with the complete redox-active moiety or with a component thereof, either before or after the probe oligonucleotide is bound to the conductive surface. The various combination possibilities of the individual steps, which lead in effect to the same test site formation unit, will be illustrated below with the aid of Figure 2 using the example of the surface hybrid $\text{Au-S(CH}_2)_2\text{-ss-oligo-spacer-UQ(RC)}$ or its more general form $\text{elec-spacer-ss-oligo-spacer-UQ(RC)}$.

[102] The reaction center can be freed of the two ubiquinone cofactors in the Q_A or Q_B binding pocket by simple manipulation (Gunner, M.R., Robertson, D.E., Dutton, P.L., 1986, Journal of Physical Chemistry, Vol. 90, pp. 3783-3795), thus yielding ubiquinone separated from the remaining RC (apoprotein including all cofactors except ubiquinone in the Q_A or Q_B binding pocket). The probe oligonucleotide has (identical or differing) reactive groups attached, via a (any) spacer, near each of its ends. In a reaction sequence "1," in the presence of a monofunctional linker (according to points a) - c) and 2.) in the section "Binding an Oligonucleotide to the Conductive Surface"), the probe oligonucleotide thus modified may be covalently attached to the electrode together with the monofunctional linker, making sure that sufficient monofunctional linker of suitable chain length is added to provide sufficient space between the individual probe oligonucleotides to permit hybridization with the target oligonucleotide and to permit the attachment of the redox-active moiety. Thereafter, UQ that was previously provided with a suitable reactive coupling group is attached to the free spacer-bridged reactive group of the probe oligonucleotide. The attachment takes place as described under a) or b) in the section "Binding a Redox-Active Moiety to a Nucleic Acid Oligomer." In the last step of this reaction sequence "1," the remaining RC (apoprotein with all cofactors except UQ) is reconstituted to UQ. In a variation hereof (reaction sequence "2"), the modified (with spacer and reactive groups) probe oligonucleotide can first be covalently bound to the

electrode without a free monofunctional linker (spacer), causing a flat attachment of the oligonucleotide. Thereafter, the free monofunctional linker (spacer) is covalently bound to the electrode. A further possibility (reaction sequence "3") is to modify the modified (with spacer and reactive groups) probe oligonucleotide first with UQ, then covalently attach it to the electrode in the presence of a free monofunctional linker (spacer), and thereafter reconstitute it with the remaining RC. Finally, in a reaction sequence "4," the modified (with spacer and reactive groups) probe oligonucleotide can first be modified with UQ so as then to reconstitute it with the remaining RC and, thereafter, to covalently bind it to the electrode. In the event that, as in the case of the RC, the redox-active moiety has a significantly greater diameter than the hybridized ds-oligonucleotide (greater than 30 Å), the covalent attachment of a suitable free monofunctional linker (spacer) to the electrode can be dispensed with; otherwise, the attachment of the structure -spacer-ss-oligo-spacer-UQ(RC) to the electrode occurs in the presence of a suitable free monofunctional linker.

[103] In the example in Figure 2, the RC, via its cofactor ubiquinone-50 (UQ) in what is known as the Q_A protein binding pocket of the RC, is covalently joined with the oligonucleotide. Alternatively, instead of the UQ cofactor in the Q_A binding pocket, another cofactor of the RC or the apoprotein can also be covalently attached to the probe oligonucleotide, any combinations of the reaction sequences "1," "2," "3," or "4" in Figure 2 may be applied, as long as they yield the same end product (cf. Figure 2), and, in any reaction steps, the probe oligonucleotide hybridized with complementary, unmodified (target) oligonucleotide may be used in place of the single-strand probe oligonucleotides. The probe oligonucleotide can also be attached directly, in other words not bridged via a spacer, to both the electrode and the redox-active moiety, as described under c) in the section "Binding a Nucleic Acid Oligomer to the Conductive Surface" or under a) in the section "Binding a Redox-Active Moiety to a Nucleic Acid Oligomer."

[104] The electrical communication between the conductive surface and the redox-active moiety bridged via a single-strand oligonucleotide in the general structure elec-spacer-ss-oligo-spacer-moiety is weak or

nonexistent. If hybridization occurs between the probe and the target, treating the test site(s) with an oligonucleotide solution to be examined leads to increased conductivity between the surface and the redox-active moiety bridged via a double-strand oligonucleotide. For the formation unit of the test site $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-UQ(RC)}$ (with 12-bp probe oligonucleotides) used as an example, this is shown schematically in Figure 3 using amperometric measurements.

[105]

The cofactor P, also called the primary donor, is electronically excited by irradiating the RC with light of a suitable wavelength, causing photoinduced charge separation within the cofactors of the RC, an electron being transferred from the excited primary donor P^* to the UQ in the Q_A binding pocket. If a suitable potential is applied to the electrode to transfer an electron from the reduced ubiquinone (UQ^-) to the electrode, current still will not flow in the case of the probe oligonucleotide not hybridized with the target oligonucleotide, since the conductivity of the ss-oligonucleotide in $\text{Au-S(CH}_2)_2\text{-ss-oligo-spacer-UQ(RC)}$ is very slight or nonexistent. In the hybridized state ($\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-UQ(RC)}$), however, conductivity is high, an electron can be transferred from UQ^- to the electrode (forming UQ), and, in the presence of a suitable redox-active substance that reduces P^* to P, the circuit is closed and further light absorption by the RC begins the cycle anew. This manifests itself amperometrically in a distinct flow of current between the electrode and the photoinducibly redox-active moiety (Figure 3). It is thus possible to detect the sequence-specific hybridization of the target with the probe oligonucleotides by amperometry using photoinduction. The individual electron transfer steps that are triggered in the surface hybrid $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-UQ(RC)}$ by light irradiation and in the presence of a suitable redox-active substance that reduces P^* to P are illustrated in Figure attachment of the RC to the probe oligonucleotide near the primary donor), the surface hybrid $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-UQ(RC)}$ can, of course, also be reversed, such that, after light irradiation, P^* is reduced by the electrode and Q^- is oxidized by a suitable oxidizing agent.

[106]

A further test site, $\text{Au-S(CH}_2)_2\text{-ss-oligo-spacer-Q-ZnBChI}$, having the general structure elec-spacer-ss-oligo-spacer-moiety, is illustrated in Figure 5. By irradiating ZnBChI with light of a suitable wavelength, ZnBChI is electronically excited and photoinduced charge separation occurs, an electron being transferred from the excited ZnBChI^* to the quinone Q. If a suitable potential is applied to the electrode to transfer to the electrode an electron from the quinone (Q^-) thus reduced, current still will not flow in the case of the probe oligonucleotide not hybridized with the target oligonucleotide, since the conductivity of the ss-oligonucleotide in $\text{Au-S(CH}_2)_2\text{-ss-oligo-spacer-Q-ZnBChI}$ is very slight or nonexistent. In the hybridized state $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-Q-ZnBChI}$, however, conductivity is high, an electron can be transferred from Q^- to the electrode (forming Q) and, in the presence of a suitable redox-active substance that reduces ZnBChI^* to ZnBChI, the circuit is closed and further light absorption by ZnBChI begins the cycle anew. This manifests itself amperometrically in a distinct flow of current between the electrode and the photoinducibly redox-active moiety (Figure 5). The sequence-specific hybridization of the target with the probe oligonucleotides can thus be detected by amperometry using photoinduction. Given suitable external conditions and a suitable attachment (e.g. $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-ZnBChI-Q}$), the surface hybrid $\text{Au-S(CH}_2)_2\text{-ds-oligo-spacer-Q-ZnBChI}$ can, of course, also be reversed, such that, after light irradiation, ZnBChI^* is reduced by the electrode and Q^- is oxidized by a suitable oxidizing agent.


[107]

Since the redox activity of the photoinducibly redox-active moiety – even at the appropriate electrode potential – is triggered only by irradiation with light of a suitable wavelength and is maintained, at most, as long as the light irradiation continues, this condition can be used to advantage, according to the present invention, in that a particular test site or group of test sites of an oligomer chip is spatially resolved by restricting the light to this test site or group of test sites. The advantage of doing so, according to the present invention, is that the various test sites (nucleic acid oligomer combinations) of an oligomer chip can be applied to a shared, continuous, electrically-conductive surface and a specific test site or a specific group of test sites can be addressed and amperometrically detected simply by

applying a suitable external potential to the (entire) surface if specifically this test site or group of test sites alone is irradiated with light. The various test sites thus need not be applied to individual (micro-) electrodes that are electrically isolated from one another and individually controllable for potential application and current selection.

[108] In addition, defective base pairings (base-pair mismatches) can be recognized by an altered cyclic voltammetric characteristic. A mismatch manifests itself in a greater potential difference between the current maximums of electroreduction and electroreoxidation (reversal of electroreduction when the potential feed direction is reversed) or electrooxidation and electrorereduction in a cyclic voltammetrically reversible electron transfer between the electrically-conducting surface and the photoinducibly redox-active moiety. This fact has an advantageous effect most importantly on amperometric detection because there, the current can be tested at a potential at which the perfectly hybridizing oligonucleotide target supplies significant current, but the defectively paired oligonucleotide target does not.

[109] **Example 1:** *Modification of ubiquinone-50 with a spacer-bridged reactive carboxylic-acid group.* The 2-methoxy group of ubiquinone-50 (UQ-50) is modified to a 2-hydroxy group by ether cleavage with HBr, a standard method (alternatively, 2-OH-UQ-50 may be produced after the method of H. W. Moore, and K. Folkers, *Journal of the American Chemical Society*, 1966, 88, pp. 564-570 or of G. Daves et al., *Journal of the American Chemical Society*, 1968, 90, pp. 5587-5593). Thereafter, in a standard method using an equimolar amount of Cl-CH₂-CH₂-CO₂H, 2-OH-UQ-50 is converted to 2-(CH₂-CH₂-CO₂H)-UQ-50 and chromatographically purified. Alternatively, in a standard method, 5-OH-6-alkyl-1,4-benzoquinone analogs of UQ-50 (produced after Catlin et al., *Journal of the American Chemical Society*, 1968, 90, pp. 3572-3574) may be modified to 5-(CH₂-CH₂-CO₂H)-UQ-50-analogs using an equimolar amount of Cl-CH₂-CH₂-CO₂H.



[110] **Example 2:** Producing the oligonucleotide electrode $Au-S(CH_2)_2$ -ss-oligo-spacerUQ(RC). $Au-S(CH_2)_2$ -ss-oligo-spacer-UQ(RC) is produced in 4 steps, namely producing the conductive surface, derivatizing the surface with the oligonucleotide probe in the presence of a suitable monofunctional linker (incubation step), covalently attaching the modified ubiquinone-50 (redox step) and reconstituting the remaining RC (reconstitution step).

[111] An approx. 100 nm thin gold film on mica (muscovite lamina) forms the support material for the covalent attachment of the double-strand oligonucleotides. For this purpose, freshly cleaved mica was purified with an argon-ion plasma in an electrical discharge chamber and gold (99.99%) was applied by electrical discharge in a layer thickness of approx. 100 nm. Thereafter, the gold film was freed of surface impurities (oxidation of

organic accumulations) with 30% H_2O_2 , / 70% H_2SO_4 and immersed in ethanol for approx. 20 minutes to dispel any oxygen adsorbed on the surface. After rinsing the surface with bidistilled water, a previously prepared 1×10^{-4} molar solution of the (modified) double-strand oligonucleotide is applied on the horizontally mounted surface, such that the entire gold surface is wetted (incubation step, see also below).

[112] For incubation, a doubly modified 12-bp single-strand oligonucleotide having the sequence 5'-TAGTCGGAAGCA-3' was used, which is esterified with $(\text{HO}-(\text{CH}_2)_2-\text{S})_2$ at the phosphate group of the 3'-end to form $\text{P-O}-(\text{CH}_2)_2-\text{S-S}-(\text{CH}_2)_2-\text{OH}$. At the 5'-end, the terminal thymine base of the oligonucleotide is modified at the C-5 carbon with $-\text{CH}=\text{CH}-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}_2$. Approximately 10^{-4} to 10^{-1} molar 2-hydroxy-mercaptoethanol (or another thiol or disulfide linker having a suitable chain length) was added to a 2×10^{-4} molar solution of this oligonucleotide in HEPES buffer (0.1 molar in water, pH 7.5 with 0.7 molar addition of TEATFB, see abbreviations) and the gold surface of a test site was completely wetted and incubated for 2-24 hours. During this reaction time, the disulfide spacer $\text{P-O}-(\text{CH}_2)_2-\text{S-S}-(\text{CH}_2)_2-\text{OH}$ of the oligonucleotide is homolytically cleaved. In this process, the spacer forms a covalent Au-S bond with Au atoms of the surface, thus causing a 1:1 coadsorption of the ss-oligonucleotide and the cleaved 2-hydroxy-mercaptoethanol. The free 2-hydroxy-mercaptoethanol that is also present in the incubation solution is likewise coadsorbed by forming an Au-S bond (incubation step).


[113] The gold electrode thus modified with a monolayer comprising ss-oligonucleotide and 2-hydroxy-mercaptoethanol was washed with bidistilled water and subsequently wetted with a solution of 3×10^{-3} molar quinone 2- $(\text{CH}_2-\text{CH}_2-\text{CO}_2\text{H})$ -UQ-50, 10^{-2} molar EDC, and 10^{-2} molar sulfo-NHS in HEPES buffer (0.1 molar (in water, pH = 7.5). After a reaction time of approx. 1 – 4 hours, the $-\text{CH}=\text{CH}-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}_2$ spacer and the 2- $(\text{CH}_2-\text{CH}_2-\text{CO}_2\text{H})$ -UQ-50 form a covalent bond (amidation between the amino group of the spacer and the C-2-acid function of 2- $(\text{CH}_2-\text{CH}_2-\text{CO}_2\text{H})$ -UQ-50, redox step).


- [114] Thereafter, the gold electrode thus modified was washed with bidistilled water and incubated for approx. 12 hours with a solution of approx. 5×10^{-5} molar ubiquinone-50-free RC in 10 mM Tris, pH = 8, with 0.7 molar addition of TEATFB at approx. 4°C, to reconstitute the remaining RC to the UQ-50 bound to the oligonucleotide (reconstitution step).
- [115] Alternatively, to covalently attach 2-(CH₂-CH₂-CO₂H)-UQ-50 to the probe oligonucleotide, under the same conditions, a 5-(CH₂-CH₂-CO₂H)-UQ-50-analog (Example 1) or another quinone of the Formula 1 – 8 having a reactive carboxylic acid may also be used, since ubiquinone-50-free RC can also be reconstituted to these.
- [116] *Example 3: Producing the oligonucleotide electrode Au-S(CH₂)₂-ss-oligo-spacer-Q-ZnBChl.* Au-S(CH₂)₂-ss-oligo-spacer-Q-ZnBChl is produced in 5 steps, namely producing the conductive surface, derivatizing the surface with the probe oligonucleotide (hybridized with a complementary strand) in the presence of a suitable monofunctional linker (incubation step), covalently attaching the electron acceptor (acceptor step), covalently attaching the electron donor (donor step), and thermally dehybridizing the double-strand oligonucleotide (dehybridization step).
- [117] The support material for the covalent attachment of the double-strand oligonucleotides, an approx. 100 nm thin gold film on mica (muscovite lamina), was produced as described in Example 1.
- [118] For incubation, a doubly modified 12-bp single-strand oligonucleotide having the sequence 5'-TAGTCGGAAGCA-3' was used, which is esterified with (HO-(CH₂)₂-S)₂ at the phosphate group of the 3'-end to form P-O-(CH₂)₂-S-S-(CH₂)₂-OH. At the 5'-end, the terminal thymine base of the oligonucleotide is modified at the C-5 carbon with -CH=CH-CO-NH-CH₂-CH₂-NH₂. A 2×10^{-4} molar solution of this oligonucleotide in the hybridization buffer (10 mM Tris, 1 mM EDTA, pH 7.5 with 0.7 molar addition of TEATFB, see abbreviations) was hybridized with a 2×10^{-4} molar solution of the (unmodified) complementary strand in the hybridization buffer at room temperature for approx. 2 hours (hybridization step). After hybridization, approx. 10^{-4} to 10^{-1} molar 2-hydroxy-mercaptoethanol (or another thiol or disulfide linker having a suitable chain length) was added to the now 1×10^{-4} molar double-strand

oligonucleotide solution and the gold surface of a test site was completely wetted and incubated for 2 – 24 hours. During this reaction time, the disulfide spacer $\text{P-O-(CH}_2\text{)}_2\text{-S-S-(CH}_2\text{)}_2\text{-OH}$ of the oligonucleotide is homolytically cleaved. In this process, the spacer forms a covalent Au-S bond with Au atoms of the surface, thus causing a 1:1 coadsorption of the ds-oligonucleotide and the cleaved 2-hydroxy-mercaptoethanol. The free 2-hydroxy-mercaptoethanol that is also present in the incubation solution is likewise coadsorbed by forming an Au-S bond (incubation step).

[119] The gold electrode thus modified with a monolayer comprising ds-oligonucleotide and 2-hydroxy-mercaptoethanol was washed with bidistilled water and subsequently wetted with a solution of 3×10^{-3} molar quinone PQQ, 10^{-2} molar EDC, and 10^{-2} molar sulfo-NHS in HEPES buffer. After a reaction time of approx. 1 – 4 hours, the $-\text{CH}=\text{CH}-\text{CO}-\text{NH}-\text{CH}_2-\text{CH}_2-\text{NH}_2$ spacer and the PQQ form a covalent bond (amidation between the amino group of the spacer and the C-7-carboxylic-acid function of the PQQ, acceptor step).

[120] Thereafter, the gold electrode thus modified was washed with bidistilled water and wetted with an aqueous solution of 3×10^{-3} molar donor ZnBChI (free acid), 1.5×10^{-1} molar EDC, 2.5×10^{-3} molar hydrazine monohydrate ($\text{NH}_2-\text{NH}_2 \cdot \text{H}_2\text{O}$), and 1×10^{-1} molar imidazole. After a reaction time of approx. 16 hours at 23°C , the C-1-carboxylic-acid function of the PQQ bound to the oligonucleotide binds via hydrazine to the free carboxylic-acid group of the ZnBChI (amidation between the amino groups of the hydrazine and the C-1-carboxylic-acid group of the PQQ or the free carboxylic-acid group of the ZnBChI, donor step). Thereafter, the double-strands were thermally dehybridized at temperatures of $T > 40^\circ\text{C}$ and rinsed again with bidistilled water (dehybridization step). The ZnBChI (free acid) is produced from Zn-BChI (produced after Hartwich et al., Journal of the American Chemical Society, 1998, 120, pp. 3684-3693) by incubation with trifluoroacetic acid.





[121] Alternatively, for example, ZnBChI (free acid) can also be bound to the 3-OH group of the 5'-terminal sugar of the probe oligonucleotide by esterification according to standard methods, or the previously covalently-joined electron-donor/electron-acceptor complex is attached to the probe oligonucleotide, as described in the donor step, via a free carboxylic-acid group, for example of the donor. In place of PQQ, under the same reaction conditions, anthraquinone-2,6-disulfonic acid disodium salt can also be used in the acceptor step. If PNA oligonucleotide is used with, for example, $\text{-NH-(CH}_2\text{)}_2\text{-N(COCH}_2\text{-base)-CH}_2\text{CO-}$ as the oligonucleotide building block, there exists an alternative possibility for attaching the ZnBChI-PQQ moiety to the nucleic acid oligomer (PNA oligonucleotide) according to d) in the section "Binding a Photoinducibly Redox-Active Moiety to a Nucleic Acid Oligomer." In this case, during PNA oligonucleotide synthesis, instead of the N-terminal base in the standard PNA synthesis reaction, PQQ is attached via its pyrrole nitrogen. Thereafter, in a manner similar to that described in the donor step, Zn-BChI is bound to the amino end of the peptide backbone by incubating the PNA oligonucleotide modified with PQQ with 3×10^{-3} molar ZnBChI (free acid), 1.5×10^{-1} molar EDC 10^{-2} and 2×10^{-1} molar sulfo-NHS in HEPES buffer (amidation between the amino group of the backbone and the carboxylic-acid group of the Zn-BChI (free acid)).